

B10

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 October 2003 (02.10.2003)

PCT

(10) International Publication Number
WO 03/079984 A2

(51) International Patent Classification⁷: A61K

(21) International Application Number: PCT/US03/08324

(22) International Filing Date: 18 March 2003 (18.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/367,045	21 March 2002 (21.03.2002)	US
60/366,915	21 March 2002 (21.03.2002)	US
60/367,048	21 March 2002 (21.03.2002)	US
60/395,468	12 July 2002 (12.07.2002)	US
60/440,305	14 January 2003 (14.01.2003)	US

(71) Applicant: PARATEK PHARMACEUTICALS, INC.
[US/US]; 75 Kneeland Street, Boston, MA 02111 (US).

(72) Inventors: NELSON, Mark, L.; 735 Worcester Road, Wellesley, MA 02481 (US). OHMENG, Kwasi; 400 Engamore Lane, Apt. 101, Norwood, MA 02062 (US). FRECHETTE, Roger; 40 Estate Lane, Reading, MA 01867 (US). ABATO, Paul; 41 Pembroke Avenue, Providence, RI 02408 (US). ASSEFA, Haregewein; 29 McCusker Drive, #5, Braintree, MA 02184 (US). BANGARAGE, Upul; 11 Bates Road, Lexington, MA 02421 (US). BERNIAC, Joel; 21 Sunrise Avenue, Stoneham, MA 02180 (US). BHATIA, Beena; 3 Brian Lane, Mansfield, MA 02048 (US). CHEN, Jackson; 1731 Beacon Street, #204, Brookline, MA 02445 (US). ISMAIL, Mohamed, Y.; 5 Hunt Road, Bedford, MA 01730 (US). KIM, Oak, A.; 86 Ellery Street, #1 Apt. 24, Cambridge, MA 02138 (US). MATHEWS, Jude; 633 Moody Street, Waltham, MA 02453 (US). MCINTYRE, Laura; 27 Thorndike Street, Arlington, MA 02474 (US). NIHLAWI, Mohammed; 230 Willard Street, #512, Quincy, MA

02169 (US). PEARSON, Andre; 23 Cutter Street, 32, Somerville, MA 02143 (US). REDDY, Laxma; 49 Eldred Street, Lexington, MA 02420 (US). SHEAHAN, Paul; 4174 Bannock Avenue, San Diego, CA 92117 (US). SIZENSKY, Emmanuelle; 4 Blackstone Lane, South Grafton, MA 01560 (US). TOURIGNY, Justin; 42 Brainerd Road, #1, Allston, MA 02134 (US). VERMA, Atul, K.; 3 Brian Lane, Mansfield, MA 02048 (US). VISKI, Peter; 887 West Roxbury Parkway, Brookline, MA 02467 (US). WARCHOL, Tadeusz; 177 South Street, Northborough, MA 01532 (US).

(74) Agents: HANLEY, Elizabeth, A. et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED TETRACYCLINE COMPOUNDS

(57) Abstract: The present invention pertains, at least in part, to novel substituted tetracycline compounds. These tetracycline compounds can be used to treat numerous tetracycline compound-responsive states, such as bacterial infections and neoplasms, as well as other known applications for minocycline and tetracycline compounds in general, such as blocking tetracycline efflux and modulation of gene expression.

WO 03/079984 A2

SUBSTITUTED TETRACYCLINE COMPOUNDS

Related Applications

This application claims priority to U.S. Provisional Patent Application
5 Serial No. 60/440,305, entitled "Substituted Tetracycline Compounds," filed January 14,
2003; U.S. Provisional Patent Application Serial No. 60/395,468, entitled "Substituted
Tetracycline Compounds," filed July 12, 2002; U.S. Provisional Patent Application
Serial No. 60/367,045, entitled "7-Substituted Tetracycline Compounds," filed March
21, 2002; U.S. Provisional Patent Application Serial No. 60/366,915, entitled "7,9-
10 Substituted Tetracycline Compounds," filed March 21, 2002; and U.S. Provisional
Patent Application Serial No. 60/367,048, entitled "9-Substituted Minocycline
Compounds," filed March 21, 2002. The entire contents of each of the aforementioned
patent applications are hereby incorporated herein by reference in their entirety.

15 Background of the Invention

The development of the tetracycline antibiotics was the direct result of a
systematic screening of soil specimens collected from many parts of the world for
evidence of microorganisms capable of producing bacteriocidal and/or bacteriostatic
compositions. The first of these novel compounds was introduced in 1948 under the
20 name chlortetracycline. Two years later, oxytetracycline became available. The
elucidation of the chemical structure of these compounds confirmed their similarity and
furnished the analytical basis for the production of a third member of this group in 1952,
tetracycline. A new family of tetracycline compounds, without the ring-attached methyl
group present in earlier tetracyclines, was prepared in 1957 and became publicly
25 available in 1967; and minocycline was in use by 1972.


Recently, research efforts have focused on developing new tetracycline
antibiotic compositions effective under varying therapeutic conditions and routes of
administration. New tetracycline analogues have also been investigated which may
prove to be equal to or more effective than the originally introduced tetracycline
30 compounds. Examples include U.S. Patent Nos. 2,980,584; 2,990,331; 3,062,717;
3,165,531; 3,454,697; 3,557,280; 3,674,859; 3,957,980; 4,018,889; 4,024,272; and
4,126,680. These patents are representative of the range of pharmaceutically active
tetracycline and tetracycline analogue compositions.

Historically, soon after their initial development and introduction, the
35 tetracyclines were found to be highly effective pharmacologically against rickettsiae; a
number of gram-positive and gram-negative bacteria; and the agents responsible for

lymphogranuloma venereum, inclusion conjunctivitis, and psittacosis. Hence, tetracyclines became known as "broad spectrum" antibiotics. With the subsequent establishment of their in vitro antimicrobial activity, effectiveness in experimental infections, and pharmacological properties, the tetracyclines as a class rapidly became widely used for therapeutic purposes. However, this widespread use of tetracyclines for both major and minor illnesses and diseases led directly to the emergence of resistance to these antibiotics even among highly susceptible bacterial species both commensal and pathogenic (e.g., pneumococci and Salmonella). The rise of tetracycline-resistant organisms has resulted in a general decline in use of tetracyclines and tetracycline analogue compositions as antibiotics of choice.

Summary of the Invention:

In one embodiment, the invention pertains, at least in part, to substituted tetracycline compounds of Formula I:



wherein:

X is $\text{CHC}(\text{R}^{13}\text{Y}^{\text{'}}\text{Y})$, CR^6R^6 , $\text{C}=\text{CR}^6\text{R}^6$, S, NR^6 , or O;

R^2 , $\text{R}^{2'}$, $\text{R}^{4'}$, and $\text{R}^{4''}$ are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R^4 is $\text{NR}^{4'}\text{R}^{4''}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;

$\text{R}^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;

R^6 and $\text{R}^{6'}$ are each independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^7 is boronic acid moiety, isoxazolyl, sulfonic acid heterocyclic, carbonyl, alkylsulfinyl, alkylsulfonyl, arylalkenyl, arylalkynyl, or $-(\text{CH}_2)_{0-3}\text{NR}^{7c}\text{C}(=\text{W}^{\text{'}})\text{WR}^{7a}$;

R^9 is hydrogen, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, thionitroso (e.g., $-N=S$), or $-(CH_2)_{0-3}NR^{9c}C(=Z')ZR^{9a}$;

Z is $CR^{9d}R^{9e}$, S, NR^{9b} or O;

5 Z' is O, S, or NR^{9f} ;

W is $CR^{7d}R^{7e}$, S, NR^{7b} or O;

W' is O, NR^{7f} S;


10 R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} , R^{9a} , R^{9b} , R^{9c} , R^{9d} , and R^{9e} are each independently hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl; and

15 Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts, esters, enantiomers, and prodrugs thereof.

20 In an embodiment, the invention pertains to 7, 9-substituted tetracycline compounds of Formula II:



wherein:

25 X is $CHC(R^{13}Y'Y)$, CR^6R^6 , S, NR^6 , or O;

R^2 , $R^{2'}$, $R^{4'}$, and $R^{4''}$ are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R^4 is $NR^{4'}R^{4''}$, alkyl, alkenyl, alkynyl, hydroxyl, halogen, or hydrogen;

30 $R^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;

R^6 and R^6' are each independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^7 is $-\text{CH}_2\text{NR}^{7a}\text{R}^{7b}$, halogen, $-\text{CH}_2\text{OR}^{7a}$, substituted alkenylfuranyl, pyrazinyl, pyridinyl, alkyl, acyl, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, or $-(\text{CH}_2)_{0-3}\text{NR}^{7c}\text{C}(=\text{W}')\text{WR}^{7a}$;

R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^9 is $-\text{CH}_2\text{SR}^{9a}$, $-\text{CH}_2\text{S}(=\text{O})\text{R}^{9a}$, $-\text{CH}_2\text{S}(=\text{O})_2\text{R}^{9a}$, $-\text{CH}_2\text{NR}^{9a}\text{R}^{9b}$, SO_3H , aminoalkyl, furanyl, substituted alkyl, $-(\text{CH}_2)_{0-3}(\text{NR}^{9c})_{0-1}\text{C}(=\text{Z}')(\text{Z})_{0-1}\text{R}^{9a}$, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, thionitroso, or $-(\text{CH}_2)_{0-3}(\text{NR}^{9c})_{0-1}\text{C}(=\text{Z}')(\text{Z})_{0-1}\text{R}^{9a}$;

Z is $\text{CR}^{9d}\text{R}^{9e}$, S, NR^{9b} or O;

Z' is O, S, or NR^{9f} ;

W is $\text{CR}^{7d}\text{R}^{7e}$, S, NR^{7b} or O;


W' is O, NR^{7f} S;

R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} , R^{9a} , R^{9b} , R^{9c} , R^{9d} , and R^{9e} are each independently hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl; and

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts, enantiomers, prodrugs, and amides thereof, provided that R^7 and R^9 are not both unsubstituted phenyl.

The invention pertains, at least in part to minocycline compounds of formula III:



5 wherein:

X is $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$, CR^6R^6 , S, NR^6 , or O;

R^2 , R^4 , $\text{R}^{4'}$, R^7 and $\text{R}^{7'}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

10 R^4 is $\text{NR}^4\text{R}^{4'}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;

R^2 , R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;

15 R^6 and R^6 are independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^7 is $\text{NR}^7\text{R}^{7'}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;

R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

20 R^9 is $-\text{C}(=\text{Z}')\text{R}^{9a}$, $\text{CH}_2\text{S}(=\text{O})\text{R}^{9a}$, $-\text{CH}_2\text{OR}^{9a}$, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, heterocyclic, arylalkenyl, arylalkynyl, thionitroso, substituted alkyl, or $-(\text{CH}_2)_{0-3}(\text{NR}^{9c})_{0-1}\text{C}(=\text{Z}')(\text{Z})_{0-1}\text{R}^{9a}$;

Z is $\text{CR}^{9d}\text{R}^{9e}$, S, NR^{9b} or O;

25 Z' is NR^{9f} , O or S;


R^{9a} , R^{9b} , R^{9c} , R^{9d} , R^{9e} and R^{9f} are each independently hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

30 R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts, amides, enantiomers, esters and prodrugs thereof.

The invention also pertains to 8 substituted tetracycline compound of the formula IV:



(IV)

wherein:

X is $CHC(R^{13}Y'Y)$, CR^6R^6 , S , NR^6 , or O ;

R^2 , R^4 , R^4' , R^7 and R^7' are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R^4 is $NR^4'R^4''$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;

R^2 , R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;

R^6 and $R^{6'}$ are independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^7 is hydrogen, hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(CH_2)_{0-3}(NR^{7c})_{0-1}C(=W')WR^{7a}$;

R^8 is hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(CH_2)_{0-3}(NR^{8c})_{0-1}C(=E')ER^{8a}$;

R^9 is hydrogen, hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(CH_2)_{0-3}NR^{9c}C(=Z')ZR^{9a}$;

- $R^{7a}, R^{7b}, R^{7c}, R^{7d}, R^{7e}, R^{7f}, R^{8a}, R^{8b}, R^{8c}, R^{8d}, R^{8e}, R^{8f}, R^{9a}, R^{9b}, R^{9c}, R^{9d}, R^{9e}$, and
 5 R^{8f} are each independently absent, hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

E is $CR^{8d}R^{8e}$, S, NR^{8b} or O;

E' is O, NR^{8f} , or S;

- 10 W is $CR^{7d}R^{7e}$, S, O or NR^{7b} ;

W' is O, NR^{7f} , or S;

R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

- Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl,
 15 amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts, esters and prodrugs thereof.

- In another further embodiment, the invention pertains, at least in part, to methods for treating subjects for tetracycline responsive states by administering to them
 20 an effective amount of a tetracycline compound of the invention, e.g., a compound of formula I, II, III, IV, or a tetracycline compound otherwise described herein.










Detailed Description of the Invention:

- The present invention pertains, at least in part, to novel substituted tetracycline
 25 compounds. These tetracycline compounds can be used to treat numerous tetracycline compound-responsive states, such as bacterial infections and neoplasms, as well as other known applications for minocycline and tetracycline compounds in general, such as blocking tetracycline efflux and modulation of gene expression.

- The term "tetracycline compound" includes many compounds with a similar ring
 30 structure to tetracycline. Examples of tetracycline compounds include: chlortetracycline, oxytetracycline, demeclocycline, methacycline, sancycline, chelocardin, rolitetracycline, lymecycline, apicycline; clomocycline, guamecycline, meglucycline, mepylcycline, penimepicycline, pipacycline, etamocycline, penimocycline, etc. Other derivatives and analogues comprising a similar four ring
 35 structure are also included (See Rogalski, "Chemical Modifications of Tetracyclines," the entire contents of which are hereby incorporated herein by reference). Table 1

depicts tetracycline and several known other tetracycline derivatives.

Table 1

 Oxytetracycline	 Demeclocycline	 Minocycline
 Methacycline	 Doxycycline	 Chlortetracycline
 Tetracycline	 Sancycline	 Chelocardin

Other tetracycline compounds which may be modified using the methods

- 5 of the invention include, but are not limited to, 6-demethyl-6-deoxy-4-dedimethylaminotetracycline; tetracyclino-pyrazole; 7-chloro-4-dedimethylaminotetracycline; 4-hydroxy-4-dedimethylaminotetracycline; 12 α -deoxy-4-dedimethylaminotetracycline; 5-hydroxy-6 α -deoxy-4-dedimethylaminotetracycline; 4-dedimethylamino-12 α -deoxyanhydrotetracycline; 7-dimethylamino-6-demethyl-6-
- 10 deoxy-4-dedimethylaminotetracycline; tetracyclinonitrile; 4-oxo-4-dedimethylaminotetracycline 4,6-hemiketal; 4-oxo-11a Cl-4-dedimethylaminotetracycline-4,6-hemiketal; 5a,6-anhydro-4-hydrazon-4-dedimethylamino tetracycline; 4-hydroxyimino-4-dedimethylamino tetracyclines; 4-hydroxyimino-4-dedimethylamino 5a,6-anhydrotetracyclines; 4-amino-4-
- 15 dedimethylamino-5a, 6 anhydrotetracycline; 4-methylamino-4-dedimethylamino tetracycline; 4-hydrazono-11a-chloro-6-deoxy-6-demethyl-6-methylene-4-dedimethylamino tetracycline; tetracycline quaternary ammonium compounds; anhydrotetracycline betaines; 4-hydroxy-6-methyl pretetramides; 4-keto tetracyclines; 5-keto tetracyclines; 5a, 11a dehydro tetracyclines; 11a Cl-6, 12 hemiketal tetracyclines;
- 20 11a Cl-6-methylene tetracyclines; 6, 13 diol tetracyclines; 6-benzylthiomethylene tetracyclines; 7, 11a -dichloro-6-fluoro-methyl-6-deoxy tetracyclines; 6-fluoro (α)-6-demethyl-6-deoxy tetracyclines; 6-fluoro (β)-6-demethyl-6-deoxy tetracyclines; 6- α -acetoxy-6-demethyl tetracyclines; 6- β acetoxy-6-demethyl tetracyclines; 7, 13-
- 25 epithiotetracyclines; oxytetracyclines; pyrazolotetracyclines; 11a halogens of tetracyclines; 12a formyl and other esters of tetracyclines; 5, 12a esters of tetracyclines;


10, 12a- diesters of tetracyclines; isotetracycline; 12-a-deoxyanhydro tetracyclines; 6-demethyl-12a-deoxy-7-chloroanhydrotetracyclines; B-nortetracyclines; 7-methoxy-6-demethyl-6-deoxytetracyclines; 6-demethyl-6-deoxy-5a-epitetracyclines; 8-hydroxy-6-demethyl-6-deoxy tetracyclines; monardene; chromocycline; 5a methyl-6-demethyl-6-deoxy tetracyclines; 6-oxa tetracyclines, and 6 thia tetracyclines.

I. 7-Substituted Tetracycline Compounds

In one embodiment, the invention pertains to novel 7-substituted tetracycline compounds.

10 The term "7-substituted tetracycline compounds" includes tetracycline compounds with substitution at the 7 position. In one embodiment, the substitution at the 7- position enhances the ability of the tetracycline compound to perform its intended function, e.g., treat tetracycline responsive states. In an embodiment, the 7-substituted tetracycline compound is 7-substituted tetracycline (e.g., wherein R^4 is $NR^{4'}R^{4''}$, $R^{4'}$ and $R^{4''}$ are methyl, R^5 is hydrogen and X is $CR^6R^{6'}$, wherein R^6 is methyl and $R^{6'}$ is hydroxy); 7-substituted doxycycline (e.g., wherein R^4 is $NR^{4'}R^{4''}$, $R^{4'}$ and $R^{4''}$ are methyl, R^5 is hydroxyl and X is $CR^6R^{6'}$, wherein R^6 is methyl and $R^{6'}$ is hydrogen); 7-substituted tetracycline compound, wherein X is $CR^6R^{6'}$, R^4 , R^5 , $R^{6'}$, and R^6 are hydrogen; or 7- substituted sancycline (wherein R^4 is $NR^{4'}R^{4''}$, $R^{4'}$ and $R^{4''}$ are methyl; R^5 is hydrogen and X is $CR^6R^{6'}$ wherein R^6 and $R^{6'}$ are hydrogen atoms.

The invention pertains, at least in part, to 7-substituted tetracycline compound of Formula I:



wherein:

X is $CHC(R^{13}Y'Y)$, $CR^6R^{6'}$, $C=CR^6R^{6'}$, S, NR^6 , or O;

R^2 , $R^{2'}$, $R^{4'}$, and $R^{4''}$ are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

30

- R^4 is $NR^{4'}R^{4''}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;
- $R^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;
- R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;
- R^6 and R^6' are each independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
- R^7 is boronic acid moiety, isoxazolyl, sulfonic acidheterocyclic, carbonyl, alkylsulfinyl, alkylsulfonyl, arylalkenyl, arylalkynyl, or $-(CH_2)_{0-3}NR^{7c}C(=W^7)WR^{7a}$;
- R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
- R^9 is hydrogen, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, thionitroso(e.g., $-N=S$), or $-(CH_2)_{0-3}NR^{9c}C(=Z')ZR^{9a}$;
- Z is $CR^{9d}R^{9e}$, S, NR^{9b} or O;
- Z' is O, S, or NR^{9f} ;
- W is $CR^{7d}R^{7e}$, S, NR^{7b} or O;
- W^7 is O, NR^{7f} S;
- R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} , R^{9a} , R^{9b} , R^{9c} , R^{9d} , and R^{9e} are each independently hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;
- R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl; and
- Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts thereof.
- In certain embodiment, R^7 is not nitro or amino.
- In an embodiment, X is $CR^6R^{6'}$; R^2 , $R^{2'}$, R^6 , $R^{6'}$, R^8 , R^9 , R^{10} , R^{11} , and R^{12} are each hydrogen; R^4 is $NR^{4'}R^{4''}$; $R^{4'}$ and $R^{4''}$ are lower alkyl (e.g., methyl); and R^5 is hydroxy or hydrogen.
- In a further embodiment, R^{7f} is hydrogen or alkyl. In another embodiment, R^{7g} is alkyl, heterocyclic, or aralkyl, e.g., benzyl.

In certain embodiments, R^{7g} and R^{7f} are linked to form a ring. R^{7g} and R^{7f} can be linked with a chain to form rings of 3-8 atoms, such as carbon (e.g., to form a piperidinyl ring), oxygen (e.g., morpholinyl, etc.), nitrogen (e.g., pyrazinyl, piperazinyl, etc.).

5 In another embodiment, R^7 is furanyl. In a further embodiment, the furan is substituted with a substituted or unsubstituted alkyl group, which itself may be substituted by one or more heterocyclic groups.

In another further embodiment, R^7 is a non-aromatic heterocycle. In a further embodiment, the heterocyclic R^7 group does not contain any double bonds, e.g. the heterocyclic R^7 group is saturated. Examples of saturated heterocyclic R^7 groups include but are not limited to, morpholine, piperazine, piperidine, etc.

In another further embodiment, R^7 comprises a carbonyl group. For example, in a further embodiment, R^7 is of the formula $-(C=O)-R^{7j}$, wherein R^{7j} is alkcarbonyl, alkyl, alkenyl, alkynyl, aryl, heterocyclic or aralkyl. In a further
15 embodiment, R^{7j} is substituted or unsubstituted alkyl. In another embodiment, the alkyl R^{7j} group is substituted with a heterocycle, such as, but not limited to morpholine, piperidine, or piperazine.

In an embodiment, R^7 is aryl. Examples of aryl R^7 groups include substituted or unsubstituted phenyl. The phenyl R^7 group can be substituted with any
20 substituent which allow the tetracycline compound to perform its intended function. Examples of substituents include, but are not limited to, alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkylloxycarbonyl, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl,
25 alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl. In certain
30 embodiments, R^7 is phenyl substituted with a sulfonyl amino or alkylcarbonylamino group.

In a further embodiment, the phenyl R^7 group is substituted with substituted or unsubstituted alkyl. Examples of substituents of the alkyl include heterocycles such as, morpholine, piperidine, and pyrrolidine. In another further
35 embodiment, the phenyl R^7 group is substituted with an amino group. The amino group also may be further substituted e.g., with an alkyl, alkenyl, alkynyl, carbonyl, alkoxy or

aryl (e.g., substituted or unsubstituted, heteroaryl, phenyl, etc.) group. The phenyl amino substituent may be substituted with any substituent or combination of substituents which allow it to perform its intended function. Examples of such substituents include halogens (e.g., fluorine, chlorine, bromine, iodine, etc.), amino (e.g., which can in turn be substituted with an alkyl, carbonyl, alkenyl, alkynyl, or aryl moiety), and arylamino (e.g., phenylamino).

The R^7 phenyl group may also be substituted with alkoxy groups. Examples of alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy, butoxy, perfluoromethoxy, perchloromethoxy, methylenedioxy, etc. The phenyl group may also be substituted with an amide group such as a carbamate moiety (e.g., an alkoxy-carbonylamino group).

The aryl group R^7 group also may be substituted or unsubstituted biaryl, e.g., naphthyl, fluorenyl, etc. The biaryl R^7 group can be substituted with any substituent which allow it to perform its intended function. Examples of substituents include but are not limited to, alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkyl-carbonyloxy, alkyloxy-carbonyl, carboxy, aryl-carbonyloxy, alkoxy-carbonylamino, alkoxy-carbonyloxy, aryloxy-carbonyloxy, carboxylate, alkyl-carbonyl, alkylamino-carbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkyl-carbonyl, aryl-carbonyl, aminoalkyl, arylalkyl-carbonyl, alkenyl-carbonyl, alkoxy-carbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfanyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

In an embodiment, the substituent is amino or formyl.

The aryl R^7 group also may be heteroaryl. Examples of heteroaryl R^7 moieties include, but are not limited to, furanyl, imidazolyl, benzothiophenyl, benzofuranyl, quinoliny, isoquinoliny, pyridinyl, pyrazolyl, benzodioxazolyl, benzoxazolyl, benzothiazolyl, benzoimidazolyl, methylenedioxyphenyl, indolyl, thienyl, pyrimidyl, pyrazinyl, purinyl, pyrazolyl, oxazolyl, isooxazolyl, naphthridinyl, thiazolyl, isothiazolyl, and deazapurinyl. In certain embodiments, the heteroaryl R^7 group is thiazolyl, thiophenyl, or furanyl.

R^7 also may be substituted or unsubstituted alkyl. The alkyl group can be a straight or branched chain, e.g., methyl, ethyl, i-propyl, n-propyl, n-butyl, i-butyl, t-butyl, pentyl, hexyl, etc. The alkyl group may also comprise a ring, e.g., a cycloalkyl (e.g., cyclopentyl, cyclohexyl, cyclopropyl, or cyclobutyl). The alkyl R^7 group may be substituted with any substituent or combination of substituents which allows the

compound to perform its intended function. Examples of substituents include, but are not limited to, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkyloxycarbonyl, carboxy, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

In certain embodiments, the alkyl group is substituted with an amino, hydroxy, carboxy, carbonyl (e.g., substituted carbonyl, e.g., morpholinyl carbonyl), heterocyclic or aryl groups. Examples heterocyclic groups include, for example, furanyl, imidazolyl, benzothiophenyl, benzofuranyl, quinolinyl, isoquinolinyl, benzodioxazolyl, benzoxazolyl, benzothiazolyl, benzoimidazolyl, methylenedioxyphenyl, indolyl, thienyl, pyridinyl, pyrazolyl, pyrimidyl, pyrazinyl, purinyl, pyrazolyl, oxazolyl, isooxazolyl, naphthridinyl, thiazolyl, isothiazolyl, and deazapurinyl. In a further embodiment, the aryl group is pyridinyl.

In a further embodiment, the aralkyl R⁷ group comprises substituted or unsubstituted phenyl. This phenyl group also may be substituted with any substituent which allows it to perform its intended function. Examples of substituents include, but are not limited to, sulfonamido, alkyl, and the other substituents listed supra for alkyl R⁷ groups.

R⁷ also may be substituted or unsubstituted alkenyl. Examples of substituents include those which allow the compound to perform its intended function. Examples of substituents include but are not limited to alkyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkyloxycarbonyl, carboxy, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl moieties.

In a further embodiment, the alkenyl R⁷ group is substituted with an aminocarbonyl (e.g., alkylaminocarbonyl, dialkylaminocarbonyl, dimethylaminocarbonyl) or alkoxycarbonyl. The alkenyl R⁷ group also may be substituted with one or more halogens (e.g., fluorine, chlorine, bromine, iodine, etc.),
5 hydroxy groups, heteroaryl groups (e.g., furanyl, imidazolyl, benzothiophenyl, benzofuranyl, quinolynyl, isoquinolynyl, benzodioxazolyl, benzoxazolyl, benzothiazolyl, benzoimidazolyl, methylenedioxyphenyl, indolyl, thienyl, pyridinyl, pyrazolyl, pyrimidyl, pyrazinyl, purinyl, pyrazolyl, oxazolyl, isooxazolyl, naphthridinyl, thiazolyl, isothiazolyl, deazapurinyl, etc.). In an embodiment, the heteroaryl substituent s
10 thiazolyl.

In a further embodiment, the alkenyl R⁷ group is substituted with a substituted or unsubstituted phenyl. The phenyl can be substituted with any substituent which allows it to perform its intended function. Examples of substituents include those listed supra for other phenyl moieties. Other examples of substituents include, but are
15 not limited to, halogens (e.g., fluorine, chlorine, bromine, iodine, etc.), alkoxy (e.g., methoxy, ethoxy, propoxy, perfluoromethyl, perchloromethyl, etc.), hydroxy, or alkyl (e.g., methyl, ethyl, propyl, butyl, pentyl, hexyl, etc.) groups.

Another example of R⁷ include substituted and unsubstituted alkynyls. The alkynyl moieties can be substituted with any substituent or combination of
20 substituents which allow the tetracycline compound of the invention to perform its intended function. Examples of the substituents include, but are not limited to alkyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkyloxycarbonyl, carboxy, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl,
25 alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl moieties.

30 In an embodiment, the alkynyl R⁷ moiety is substituted with an aryl, e.g., substituted or unsubstituted heteroaryl, phenyl, etc. This aryl moiety may be substituted with any substituent or combinations of substituents listed supra for the alkynyl R⁷ moiety. Examples of advantageous substituents include, but are not limited to, carbonylamino (e.g., alkylcarbonylamino, dialkylcarbonylamino, arylcarbonylamino,
35 etc.) and sulphonamido groups.

In another embodiment, the alkynyl R^7 group is substituted with a tetracycline moiety. The term "tetracycline moiety" includes a four ring tetracycline ring system as described above. This may be connected to the alkynyl R^7 group through a linker of 1-20 atoms. The linker may be attached to the tetracycline moiety at any position on that ring system which is convenient or allows the compound to perform its intended function. In a certain embodiment, the tetracycline moiety is attached to the linker at its 7 position.

Other examples of R^7 moieties include substituted and unsubstituted alkylcarbonyl amino, sulfonamido, imino and carbonyl moieties. The carbonyl moieties may be substituted with a substituted or unsubstituted alkyl group. Examples of possible substituents of the alkyl group include, but are not limited to, aryl moieties such as phenyl and heteroaryls (e.g., pyridinyl, etc.). Examples of substituents of the imino group include, but are not limited to, hydroxy and alkoxy groups.

In another embodiment, R^7 is $NR^{7c}(C=W')WR^{7a}$. Examples of tetracycline compounds of the invention include compounds wherein R^{7c} is hydrogen, W' is oxygen and W is oxygen. In certain embodiments, R^{7a} is substituted or unsubstituted phenyl. Examples of substituents include, but are not limited to, alkyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkyloxycarbonyl, carboxy, alkylcarbonylamino, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl. In a further embodiment, R^{7a} is substituted or unsubstituted alkyl.

In a further embodiment, R^4 is $NR^{4'}R^{4''}$, X is CR^6R^6' ; R^2 , R^2' , R^6 , R^6' , R^8 , R^9 , R^{10} , R^{11} , and R^{12} are each hydrogen; $R^{4'}$ and $R^{4''}$ are lower alkyl; and R^5 is hydroxy or hydrogen. $R^{4'}$ and $R^{4''}$ may be methyl, for example, and R^5 may be hydrogen.

In a further embodiment, R^7 is a substituted pyrrole. The pyrrole may be substituted with a carbonyl moiety, such as, for example substituted or unsubstituted alkoxycarboynyl.

In another embodiment, R^7 is a substituted pyrazine. Examples of substituents include substituted or unsubstituted alkoxycarbonylamino groups, an alkaminocarbonylamino groups, and alkylaminocarbonyl groups.

In another embodiment, R^7 is substituted pyridine. Examples of substituents include halogens. In another embodiment, R^7 is alkenyl and substituted with a halogen (e.g., fluorine), or a carbonyl group. In another embodiment R^7 is pyrazolyl.

- 5 In another embodiment, R^7 is $-C(=W')WR^{7a}$, and W' is NR^{7f} , W is $CR^{7d}R^{7e}$, and R^{7a} is hydrogen. In a further embodiment, R^{7f} is alkoxy.


In yet another embodiment, R^7 is $-NR^{7c}C(=W')WR^{7a}$, R^{7c} is hydrogen, W' is NH , W is NR^{7b} , and R^{7a} and R^{7b} taken together are heterocyclic, e.g., pyrrolidine.


- 10 In another embodiment, R^7 is a boronic acid. Examples of boronic acids include compounds wherein R^7 is $-B(OR^{b1})(OR^{b2})$ and R^{b1} and R^{b2} are each hydrogen or alkyl, and optionally linked to form a ring.


In another embodiment, R^7 is a sulfonic acid, or an ester or prodrug thereof. Examples of sulfonic acids include $-SO_3H$.


- 15 In yet another embodiment, R^7 is isoxazolyl. R^7 may be substituted or unsubstituted. In one embodiment, the isooxazole is substituted with, for example, alkyl groups.


Other examples of tetracycline compounds include the following:






































































and pharmaceutically acceptable enantiomers, amides, salts, prodrugs, and esters thereof.

5 2. 7,9-Substituted Tetracycline Compounds

The invention also pertains, at least in part to 7,9-substituted tetracycline compounds.

The term "7,9-substituted tetracycline compounds" includes tetracycline compounds with substitution at the 7 and 9- positions. In one embodiment, the substitution at the 7- and 9- positions enhances the ability of the tetracycline compound to perform its intended function, e.g., treat tetracycline responsive states. In an embodiment, the 7,9-substituted tetracycline compound is 7,9-substituted tetracycline (e.g., wherein R^4 is $NR^{4'}R^{4''}$; $R^{4'}$ and $R^{4''}$ are methyl, R^5 is hydrogen and X is CR^6R^6 , wherein R^6 is methyl and R^6 is hydroxy); 7,9-substituted doxycycline (e.g., wherein R^4 is $NR^{4'}R^{4''}$; $R^{4'}$ and $R^{4''}$ are methyl, R^5 is hydroxyl and X is CR^6R^6 , wherein R^6 is methyl and R^6 is hydrogen); or 7, 9- substituted sancycline (wherein R^4 is $NR^{4'}R^{4''}$; $R^{4'}$ and $R^{4''}$ are methyl; R^5 is hydrogen and X is CR^6R^6 wherein R^6 and R^6 are hydrogen atoms. In an embodiment, the substitution at the 7 position of the 7, 9-substituted tetracycline compound is not chlorine or trimethylamino. In one embodiment, R^4 is hydrogen.

20 The 7, 9-substituted tetracycline compounds of the invention include compounds of Formula II:



wherein:

25 X is $CHC(R^{13}Y'Y)$, CR^6R^6 , S, NR^6 , or O;
 R^2 , R^2 , $R^{4'}$, and $R^{4''}$ are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

- R^4 is $NR^{4'}R^{4''}$, alkyl, alkenyl, alkynyl, hydroxyl, halogen, or hydrogen;
 R^2 , R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;
 R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl,
 heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl,
 5 alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;
 R^6 and R^6 are each independently hydrogen, methylene, absent,
 hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl,
 alkylsulfonyl, alkylamino, or an arylalkyl;
 R^7 is $-CH_2NR^{7a}R^{7b}$, halogen, $-CH_2OR^{7a}$, substituted alkenylfuranyl,
 10 pyrazinyl, pyridinyl, alkyl, acyl, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio,
 alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, or $-(CH_2)_{0-3}NR^{7c}C(=W')WR^{7a}$;
 R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl,
 alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
 15 R^9 is $-CH_2SR^{9a}$, $-CH_2S(=O)R^{9a}$, $-CH_2S(=O)_2R^{9a}$, $-CH_2NR^{9a}R^{9b}$, SO_3H ,
 aminoalkyl, furanyl, substituted alkyl, $-(CH_2)_{0-3}(NR^{9c})_{0-1}C(=Z')(Z)_{0-1}R^{9a}$, nitro, alkyl,
 alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino,
 arylalkenyl, arylalkynyl, thionitroso, or $-(CH_2)_{0-3}(NR^{9c})_{0-1}C(=Z')(Z)_{0-1}R^{9a}$;
 Z is $CR^{9d}R^{9c}$, S, NR^{9b} or O;
 20 Z' is O, S, or NR^{9f} ;
 W is $CR^{7d}R^{7e}$, S, NR^{7b} or O;
 W' is O, NR^{7f} S;
 R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} , R^{9a} , R^{9b} , R^{9c} , R^{9d} , and R^{9e} are each independently
 hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl,
 25 alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;
 R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio,
 alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl; and
 Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano,
 sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl,
 30 alkylamino, or an arylalkyl, and pharmaceutically acceptable salts thereof, provided that
 R^7 and R^9 are not both unsubstituted phenyl.

The tetracycline compounds of the invention include, for example,
 compounds wherein X is $CR^6R^{6'}$; R^4 is $NR^{4'}R^{4''}$; R^2 , $R^{2'}$, R^6 , $R^{6'}$, R^8 , R^{10} , R^{11} , and R^{12}
 are each hydrogen; $R^{4'}$ and $R^{4''}$ are lower alkyl; and R^5 is hydroxy or hydrogen. In an
 35 embodiment, $R^{4'}$ and $R^{4''}$ are each methyl and R^5 is hydrogen. The tetracycline

compounds of the invention include each possible combination of R^7 and R^9 substituents discussed below.

In an embodiment, R^7 is alkylamino, e.g., $CH_2-NR^{7g}R^{7f}$. In a further embodiment, R^{7f} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heterocyclic, or aralkyl. R^{7f} may be substituted with any substituent which allows the tetracycline compound to perform its intended function. In a further embodiment, R^{7g} is alkyl, alkenyl, alkynyl, aryl, heterocyclic, or hydrogen.

In another embodiment, R^7 may be substituted or unsubstituted aryl, e.g., heteroaryl, e.g., furanyl. In another embodiment, R^7 may be alkyl, e.g., methyl, ethyl, propyl, etc.

In an embodiment, R^7 is aryl (e.g., heteroaryl or substituted or unsubstituted phenyl). The phenyl R^7 group may be substituted with one or more substituents. Examples of substituents of phenyl R^7 groups include alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkylloxycarbonyl, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl. In certain embodiments, the substituent is substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, butyl, etc.), nitro, halogen (e.g., fluorine, bromine, chlorine, iodine, etc.), amino (e.g., unsubstituted amino, alkyl amino, dialkylamino (e.g., dimethylamino), or alkoxy (methylenedioxy or methoxy).

R^7 also may be substituted or unsubstituted alkyl (e.g., methyl, ethyl, i-propyl, n-propyl, t-butyl, i-butyl, n-butyl, pentyl, n-pentyl, n-hexyl, or hexyl). The alkyl may be branched or straight chain and may comprise a ring, e.g., a cycloalkyl ring, e.g., cyclohexyl ring.

The alkyl R^7 group may be substituted with any substituent which allows the tetracycline compound to perform its intended function. Examples of substituents include, but are not limited to, alkenyl, halogen (e.g., fluorine, chlorine, bromine, iodine, etc.), hydroxyl, alkoxy (e.g., methoxy, ethoxy, perchloromethoxy, perfluoromethoxy, etc.), alkylcarbonyloxy, alkylloxycarbonyl, arylcarbonyloxy, arylcarbonylamino, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl,

alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

Examples of heterocyclic substituents include N-isoindeole-[1,3]-dione (e.g., phthalimide). In an embodiment, the substituent is arylcarbonylamino, e.g., heteroaryl carbonyl amino. The heteroaryl group may be, for example, pyridinyl. Other examples of substituents include amino or carboxylate.

In another embodiment, R^7 is acyl, e.g., acetyl.

In yet another embodiment, R^7 is substituted or unsubstituted alkynyl.

Examples of substituents include those which allow the tetracycline compound to perform its intended function. Examples of substituents include, but are not limited to, alkyl (e.g., methyl, ethyl, propyl, butyl, etc.), alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkylloxycarbonyl, arylcarbonyloxy, arylcarbonylamino, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

In a further embodiment, the aryl substituent is substituted or unsubstituted phenyl. The phenyl also may be further substituted with one or more substituents which allow the compound to perform its intended function. Examples of phenyl substituents include, but are not limited to, alkoxy (e.g., methoxy).

Any of the above described R^7 groups may be combined with any of the groups described below as R^9 groups. In an embodiment, the invention pertains to tetracycline compounds wherein R^9 is aminoalkyl. Examples of aminoalkyl R^9 groups include groups of the formula: $-\text{CH}_2-\text{NR}^{9g}\text{R}^{9f}$. Examples of R^{9f} groups include substituted and unsubstituted alkyl, alkenyl, alkynyl, aryl, heterocyclic, aralkyl, and hydrogen. In a further embodiment, R^{9f} may be further substituted with any substituent which allow s the tetracycline compound to perform its intended function, for example, treat tetracycline associated states. Examples of R^{9g} groups include substituted and unsubstituted alkyl, alkenyl, alkynyl, aryl, heterocyclic, aralkyl, and hydrogen. In a further embodiment, R^{9g} may be further substituted with any substituent which allow s

the tetracycline compound to perform its intended function, for example, treat tetracycline associated states.

In a further embodiment, R^9 is thioalkyl, e.g., $-\text{CH}_2\text{-S-}R^{9h}$. In a further embodiment, R^{9h} is hydrogen, alkyl, alkenyl, alkynyl, heterocyclic, or aralkyl. In one embodiment, R^{9h} is alkyl.

The tetracycline compounds of the invention include compounds wherein R^9 is substituted or unsubstituted aryl (e.g., carbocyclic or heteroaryl). In an embodiment, R^9 is substituted or unsubstituted phenyl. The substituted phenyl group can be substituted with any substituent or combination of substituents which allows the compound to perform its intended function. Examples of substituents include, but are not limited to, alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkyloxy carbonyl, arylcarbonyloxy, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

In a further embodiment, the phenyl R^9 substituent is substituted or unsubstituted alkyl, nitro, halogen, amino, or alkoxy (e.g., methylenedioxy).

The invention also includes compounds wherein R^9 is substituted or unsubstituted alkyl (e.g., methyl, ethyl, i-propyl, n-propyl, i-butyl, t-butyl, n-butyl, pentyl, hexyl, etc.). The alkyl group may be substituted with any substituent that allows the compound to perform its intended function. Examples of the substituents include, but are not limited to, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkyloxy carbonyl, arylcarbonyloxy, arylcarbonylamino, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxy carbonyloxy, alkylcarbonylamino, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

In an embodiment, wherein said substituent is unsubstituted amino. In a further embodiment, the R^9 group is aminomethyl. In another, the alkyl R^9 group is substituted with arylcarbonylamino (e.g., heteroarylcarbonylamino, e.g., pyridinylcarbonylamino) or alkylcarbonylamino.

5 In another further embodiment, the R^9 alkyl group is substituted with a heterocyclic substituent, such as isoindole-[1,3]-dione (e.g., phthalimide).

In an embodiment, R^7 is acyl, e.g., acetyl.

In yet another embodiment, R^9 is substituted or unsubstituted alkynyl. The alkynyl R^9 group can be substituted with any substituent which allows the
 10 tetracycline compound of the invention to perform its intended function. Examples of substituents include, but are not limited to, alkyl (e.g., methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, etc.), alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkylloxycarbonyl, arylcarbonyloxy, arylcarbonylamino, alkoxycarbonylamino, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl,
 15 alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl,
 20 alkylaryl, aryl and heteroaryl.

In a further embodiment, the aryl substituted alkynyl R^9 moiety is, for example, substituted or unsubstituted phenyl. The phenyl may be substituted with, for example, alkoxy, e.g., methoxy. Examples of alkenyl substituents include cycloalkenes such as, cyclohexene.

25 In one embodiment, R^9 is not unsubstituted phenyl when R^7 is unsubstituted phenyl.

In one embodiment, R^7 is pyridinyl and R^9 is alkyl substituted with piperidine or alkylamino. In a further embodiment, R^9 is $-\text{CH}_2-\text{S}(=\text{O})_2\text{-alkyl}$, $-\text{SO}_3\text{H}$, iodo, $\text{NO}_2\text{-CH}_2\text{S-alkyl}$, $-\text{CH}_2-\text{S}(=\text{O})\text{-alkyl}$, amino, nitro, or $-\text{CH}_2\text{O-alkyl}$.

30 In another embodiment, R^7 is furanyl, and R^9 is substituted or unsubstituted aminoalkyl, or alkyl substituted with substituted or unsubstituted morpholine or piperidine.


In yet another embodiment R^7 is $-\text{C}(=\text{W}')\text{WR}^{7a}$, wherein W' is NR^{7f} , W is $\text{CR}^{7d}\text{R}^{7e}$, and R^{7a} is hydrogen and R^9 is aminoalkyl. Examples of R^{7f} include alkoxy
 35 and examples of R^9 include alkylaminoalkyl. Each of these substituents may be further substituted.


In another embodiment, R^7 is iodo or chloro. In a further embodiment, R^9 is .

In another embodiment, R^7 is $-\text{CH}_2\text{O-alkyl}$ and R^9 is alkyl.

Examples of 7,9-substituted tetracycline compounds of the invention

5 include those listed below:





and pharmaceutically acceptable salts, esters, enantiomers, prodrugs, and amides thereof.


3. 9-Substituted Tetracycline Compounds

5 In another embodiment, the invention pertains to 9-substituted tetracycline compounds.

The term "9-substituted tetracycline compounds" includes tetracycline compounds with substitution at the 9 position. In one embodiment, the substitution at the 9- position enhances the ability of the tetracycline compound to perform its intended
 10 function, e.g., treat tetracycline responsive states. In an embodiment, the 9-substituted tetracycline compound is 9-substituted tetracycline (e.g., wherein R^4 is $NR^{4'}R^{4''}$, $R^{4'}$ and $R^{4''}$ are methyl, R^5 is hydrogen and X is $CR^6R^{6'}$, wherein R^6 is methyl and $R^{6'}$ is hydroxy, and R^7 is hydrogen); 9-substituted doxycycline (e.g., wherein R^4 is $NR^{4'}R^{4''}$, $R^{4'}$ and $R^{4''}$ are methyl, R^5 is hydroxyl and X is $CR^6R^{6'}$, wherein R^6 is methyl and $R^{6'}$ is
 15 hydrogen, and R^7 is hydrogen); 9- substituted minocycline (wherein R^4 is $NR^{4'}R^{4''}$, $R^{4'}$ and $R^{4''}$ are methyl; R^5 is hydrogen and X is $CR^6R^{6'}$ wherein R^6 and $R^{6'}$ are hydrogen atoms, and R^7 is dimethylamino); 9-substituted 4-dedimethylamino tetracycline compound, wherein X is $CR^6R^{6'}$, R^4 , R^5 , R^6 , $R^{6'}$, and R^7 are hydrogen; and 9- substituted sancycline (wherein R^4 is $NR^{4'}R^{4''}$, $R^{4'}$ and $R^{4''}$ are methyl; R^5 and R^7 are hydrogen and
 20 X is $CR^6R^{6'}$ wherein R^6 and $R^{6'}$ are hydrogen atoms).

The invention also pertains, at least in part, to tetracycline compounds of

Formula III:



wherein:

- 5 X is $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$, $\text{CR}^{6'}\text{R}^6$, S, NR^6 , or O;
 R^2 , $\text{R}^{4'}$, $\text{R}^{4''}$, R^7 and $\text{R}^{7'}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;
 R^4 is $\text{NR}^{4'}\text{R}^{4''}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;
- 10 $\text{R}^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;
 R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;
 R^6 and $\text{R}^{6'}$ are independently hydrogen, methylene, absent, hydroxyl, halogen,
- 15 thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
 R^7 is $\text{NR}^{7'}\text{R}^{7''}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;
 R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
- 20 R^9 is $-\text{C}(=\text{Z}')\text{R}^{9a}$, $\text{CH}_2\text{S}(=\text{O})\text{R}^{9a}$, $-\text{CH}_2\text{OR}^{9a}$, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, heterocyclic, arylalkenyl, arylalkynyl, thionitroso, substituted alkyl, or $-(\text{CH}_2)_{0-3}(\text{NR}^{9c})_{0-1}\text{C}(=\text{Z}')(\text{Z})_{0-1}\text{R}^{9a}$;
 Z is $\text{CR}^{9d}\text{R}^{9e}$, S, NR^{9b} or O;
 Z' is NR^{9f} , O or S;
- 25 R^{9a} , R^{9b} , R^{9c} , R^{9d} , R^{9e} and R^{9f} are each independently hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;
 R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
- 30 R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts, esters and prodrugs thereof.

5

In an embodiment, the invention features compounds wherein X is CR^6R^6' ; $R^2, R^{2'}, R^5, R^6, R^{6'}, R^8, R^9, R^{10}, R^{11}$, and R^{12} are each hydrogen; R^4 is $NR^{4'}R^{4''}$; R^7 is $NR^{7'}R^{7''}$ and $R^{4'}, R^{4''}, R^{7'}$, and $R^{7''}$ are each lower alkyl, e.g., methyl. In certain other embodiments, R^7 may be hydrogen.

10

In one embodiment, R^9 is alkyl amino. For example, R^9 may be $-CH_2-NR^{9f}R^{9g}$. Examples of R^{9f} include but are not limited to hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, etc. R^{9f} may be substituted with any substituent which allows the minocycline compound of the invention to perform its intended function, e.g., treat tetracycline associated states or other functions. In a further embodiment, R^{9f} is

15 hydrogen or alkyl.

In other embodiments, R^{9f} and R^{9g} are linked by a chain of from 0 to 5 atoms. The atoms may be carbon, oxygen, nitrogen, sulfur, etc. In certain embodiments, the ring that is formed has 3, 4, 5, 6, or 7 members. In other embodiments, the ring that is formed is morpholinyl, piperidinyl, or pyrazinyl. The ring

20 may be substituted or unsubstituted.

In certain embodiments, R^{9g} is heterocyclic (e.g., piperidinyl, piperazinyl, morpholinyl, pyrazolyl, pyridinyl, etc.) In other embodiments, R^{9g} is aralkyl. R^{9g} may comprise a substituted or unsubstituted heteroaryl group, or may comprise a substituted or unsubstituted phenyl group. In other embodiments, R^{9g} is substituted or unsubstituted

25 heteroaryl or phenyl.

In other embodiments, R^9 is substituted amino. In certain embodiments, R^9 is amino substituted with a heterocyclic group, such as, but not limited to, morpholinyl, piperazinyl, or piperidinyl. In other embodiments, R^9 is substituted or unsubstituted aralkylamino, e.g., substituted or unsubstituted benzylamino.

30

In yet other embodiments, R^9 may also be furanyl. In further embodiments, furanyl R^9 groups may be substituted with one or more substituents, such as, but not limited to, heterocyclic groups and other moieties which allow the minocycline compound of the invention to perform its intended function.

Other examples of R^9 include substituted and unsubstituted aryl groups.

35

The aryl groups include substituted and unsubstituted heteroaryls (e.g., furanyl, imidazolyl, benzothiophenyl, benzofuranyl, quinolinyl, isoquinolinyl, benzodioxazolyl,

benzoxazolyl, benzothiazolyl, benzoimidazolyl, methylenedioxyphenyl, indolyl, thienyl, pyrimidyl, pyrazinyl, purinyl, pyrazolyl, oxazolyl, isooxazolyl, naphthridinyl, thiazolyl, isothiazolyl, or deazapurinyl), substituted or unsubstituted phenyl, and groups with more than one aromatic ring, such as naphthyl.

- 5 Examples of substituents of R⁹ include, but are not limited to, alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkylloxycarbonyl, arylcarbonyloxy, alkoxy carbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxy carbonyl, silyl, aminocarbonyl, 10 alkylthiocarbonyl, phosphate, aralkyl, phosphonate, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

- 15 In a further embodiment, the aryl R⁹ group is substituted with one or more substituents such as, for example, carboxylate, alkyl, alkenyl, alkynyl, aryl, heterocyclic, cyano, amino, halogen, alkoxy, alkoxy carbonyl, amido, alkylcarbonyl, or nitro.

- 20 In another embodiment, R⁹ is substituted or unsubstituted alkynyl. The alkynyl R⁹ group may be substituted with a substituted or unsubstituted aryl group, such as, for example, phenyl. The possible substituents for the substituted phenyl group include, for example, those listed *supra*, for the aryl R⁹ group. Furthermore, the substituted alkynyl R⁹ group may be substituted with a heteroaryl (e.g., pyridinyl), alkyl (e.g., methyl, ethyl, propyl, butyl, pentyl, hexyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, *etc.*), alkenyl (e.g., ethenyl, propenyl, butenyl, pentenyl, hexenyl, *etc.*), 25 carboxylate, silyl (e.g., trialkylsilyl, e.g., trimethylsilyl), aralkyl, or an alkylloxycarbonyl group.

- Each of these groups may also be further substituted, with such substituents as alkyl, alkenyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, alkylloxycarbonyl, arylcarbonyloxy, alkoxy carbonyloxy, aryloxycarbonyloxy, 30 carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aminoalkyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxy carbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, aralkyl, phosphonate, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, sulfate, arylthio, thiocarboxylate, alkylsulfinyl, sulfonate, sulfamoyl, 35 sulfonamido, nitro, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

In a further embodiment, the alkynyl R^9 group is substituted with an aminoalkyl group. The aminoalkyl group may then also be substituted with, for example, an alkyl, alkenyl, alkynyl, acyl, carbonyl, or alkylsulfone group.

In another further embodiment, the alkynyl R^9 group is substituted with a
5 cycloalkenyl group, such as, for example, cyclopentene.

In another embodiment, R^9 is alkyl. The alkyl group may be substituted or unsubstituted. Examples of alkyl groups include, for example, both straight chain, branched and cyclic alkyl groups. For example, alkyl groups include methyl, ethyl, i-propyl, n-propyl, i-butyl, n-butyl, t-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, *etc.*
10 Cyclic alkyl groups include groups with one or more rings, such as, for example, cyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, *etc.* In an embodiment, the alkyl R^9 group is 2-cyclopentylethyl.

Examples of substituents of alkyl groups include, for example, halogens (*e.g.*, fluorine, chlorine, bromine, iodine, *etc.*), hydroxyl, alkoxy (*e.g.*, methoxy, ethoxy,
15 propoxy, butoxy, pentoxy, perfluoromethoxy, perchloromethoxy, *etc.*), alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, carboxy, alkylcarbonyl, arylcarbonyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate,
20 phosphonate, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfate, alkylsulfinyl, alkenyl, sulfonate, sulfamoyl, sulfonamido, nitro, alkenyl, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl.

In another embodiment, the minocycline compound of the invention is a compound wherein R^9 is $-NR^{9c}C(=Z')ZR^{9a}$, $-CH_2NR^{9c}C(=Z')ZR^{9a}$, $-(CH_2)_2NR^{9c}C(=Z')ZR^{9a}$, or $-(CH_2)_3NR^{9c}C(=Z')ZR^{9a}$. In certain embodiments, R^9 is $-NR^{9c}C(=Z')ZR^{9a}$ or $-CH_2NR^{9c}C(=Z')ZR^{9a}$. Examples of R^{9c} include hydrogen. Z' may be, for example, S, NH, or O. Examples of Z include NR^{9b} (*e.g.*, when R^{9b} is hydrogen, alkyl, *etc.*), O or S.
25

Examples of R^{9a} groups include aryl groups such as substituted and
30 unsubstituted phenyl. Examples of possible substituents of aryl R^{9a} groups include, but are not limited to, alkyl (*e.g.*, methyl, ethyl, propyl, butyl, pentyl, hexyl, perfluoromethyl, perchloroethyl, *etc.*), alkenyl, halogen (*e.g.*, fluorine, chlorine, bromine, iodine, *etc.*), hydroxyl, alkoxy (*e.g.*, methoxy, ethoxy, propoxy, perfluoromethoxy, perchloromethoxy, *etc.*), alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy,
35 aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, arylalkylcarbonyl,

- alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonato, phosphinato, cyano, amino, acylamino, amido, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, acetyl, alkyl, cyano, azido, heterocyclyl, alkylaryl, aryl and heteroaryl groups.

- In certain embodiments, at least one of the substituents of the substituted phenyl is nitro, alkoxy (*e.g.*, methoxy, methylenedioxy, perfluoromethoxy) alkyl (*e.g.*, methyl, ethyl, propyl, butyl, or pentyl), acetyl, halogen (*e.g.*, fluorine, chlorine, bromine, or iodine), or amino (*e.g.*, dialkylamino). In certain embodiments, the alkoxy group is perhalogenated, *e.g.*, perfluoromethoxy.

- Examples of aryl R^{9a} groups include, but are not limited to, unsubstituted phenyl, *para*-nitrophenyl, *para*-methoxy phenyl, *para*-perfluoromethoxy phenyl, *para*-acetyl phenyl, 3, 5-methylenedioxyphenyl, 3,5-diperfluoromethyl phenyl, *para*-bromo phenyl, *para*-chloro phenyl, and *para*-fluoro phenyl.

- Other examples of aryl R^{9a} groups include substituted and unsubstituted heterocycles (*e.g.*, furanyl, imidazolyl, benzothiophenyl, benzofuranyl, quinolinyl, isoquinolinyl, benzodioxazolyl, benzoxazolyl, benzothiazolyl, benzoimidazolyl, methylenedioxyphenyl, indolyl, thienyl, pyrimidyl, pyrazinyl, purinyl, pyrazolyl, pyrrolidinyl, oxazolyl, isooxazolyl, naphthridinyl, thiazolyl, isothiazolyl, or deazapurinyl) and substituted and unsubstituted biaryl groups, such as naphthyl and fluorene.

- R^{9a} also may be substituted or unsubstituted alkyl, *e.g.*, methyl, ethyl, propyl, butyl, pentyl, *etc.* Examples of substituents include but are not limited to halogens (*e.g.*, fluorine, bromine, chlorine, iodine, *etc.*), hydroxyl, alkoxy (*e.g.*, methoxy, ethoxy, propoxy, butoxy, *etc.*), alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy, carboxylate, alkylcarbonyl, alkylamino, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, silyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonato, phosphinato, cyano, amino, acylamino, amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfate, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, alkenyl, heterocyclyl, alkylaryl, aryl and heteroaryl.

- R^{9a} also can be substituted or unsubstituted alkenyl. Examples of substituents for alkenyl R^{9a} groups include those listed above for alkyl R^{9a} groups. Examples of alkenyl R^{9a} groups include pent-1-enyl.


In an embodiment, Z' is NH, Z is NH, and R^{9a} is alkyl.


In another embodiment, R^9 is alkyl and substituted with a heterocycle, such as 2,3-dihydro-isoindole.


In another embodiment, R^9 is $-C(=Z')R^{9a}$, Z' is NR^{9f} , and R^{9a} is hydrogen. R^{9f} may be alkoxy.


In a further embodiment, R^9 is substituted aminoalkyl. R^9 may be substituted, for example, with a substituted or unsubstituted alkyloxycarbonyl group, a substituted or unsubstituted alkyl group and/or a substituted or unsubstituted aralkyl group.

10 In another embodiment, the invention pertains to tetracycline compounds of the formulae:










or pharmaceutically acceptable salts, amides, enantiomers, prodrugs, or esters thereof.

4. 8-Substituted Tetracycline Compounds

5 The invention also pertains, at least in part to 8-substituted tetracycline compounds.

The term "8-substituted tetracycline compounds" includes tetracycline compounds with substitution at the 8- position. In one embodiment, the substitution at the 8- position enhances the ability of the tetracycline compound to perform its intended function, e.g., treat tetracycline responsive states. In an embodiment, the 8-substituted tetracycline compound is 8-substituted tetracycline (e.g., wherein R^4 is $NR^{4'}R^{4''}$; $R^{4'}$ and $R^{4''}$ are methyl, R^5 is hydrogen and X is CR^6R^6 , wherein R^6 is methyl and R^6 is hydroxy); 8-substituted doxycycline (e.g., wherein R^4 is $NR^{4'}R^{4''}$; $R^{4'}$ and $R^{4''}$ are methyl, R^5 is hydroxyl and X is CR^6R^6 , wherein R^6 is methyl and R^6 is hydrogen); or 15 8- substituted sancycline (wherein R^4 is $NR^{4'}R^{4''}$; $R^{4'}$ and $R^{4''}$ are methyl; R^5 is hydrogen and X is CR^6R^6 wherein R^6 and R^6 are hydrogen atoms. In an embodiment, the substitution at the 7 position of the 8-substituted tetracycline compound is not chlorine or trimethylamino. In one embodiment, R^4 is hydrogen.

20 In one embodiment, the 8-substituted tetracycline compound is of formula IV:



wherein:

25 X is $CHC(R^{13}Y'Y)$, CR^6R^6 , S, NR^6 , or O;
 R^2 , $R^{4'}$, $R^{4''}$, R^7 and R^7 are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

- R^4 is $NR^{4'}R^{4''}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;
 R^2 , R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;
 R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, 5 arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;
 R^6 and R^6' are independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
 R^7 is hydrogen, hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, 10 alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(CH_2)_{0-3}(NR^{7c})_{0-1}C(=W')WR^{7a}$;
 R^8 is hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(CH_2)_{0-3}(NR^{8c})_{0-1}C(=E')ER^{8a}$;
 R^9 is hydrogen, hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, 15 alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(CH_2)_{0-3}NR^{9c}C(=Z')ZR^{9a}$;
 R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} , R^{7f} , R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} , R^{8f} , R^{9a} , R^{9b} , R^{9c} , R^{9d} , R^{9e} , and R^{8f} are each independently absent, hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, 20 alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;
 E is $CR^{8d}R^{8e}$, S, NR^{8b} or O;
 E' is O, NR^{8f} , or S;
 W is $CR^{7d}R^{7e}$, S, O or NR^{7b} ;
 W' is O, NR^{7f} , or S;
 R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
 Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, 30 alkylamino, or an arylalkyl, and pharmaceutically acceptable salts, esters and prodrugs thereof.

In a further embodiment, the invention pertains to compounds wherein X is CR^6R^6' ; R^2 , R^2' , R^6 , R^6' , R^8 , R^{10} , R^{11} , and R^{12} are each hydrogen; R^4 is $NR^{4'}R^{4''}$; $R^{4'}$ and $R^{4''}$ are lower alkyl; and R^5 is hydroxy or hydrogen.


In another embodiment, R^8 is halogen, e.g., chloro or bromo.

In another embodiment, R^8 is unsubstituted or substituted aryl. Examples of heteroaryl group include furanyl. Examples of aryl groups include phenyl.

In a further embodiment, R^9 or R^7 is a substituted or unsubstituted amino.

- 5 The amino group may be substituted, for example, with carbonyl, alkyl, or any other substituent described herein.


Examples of 8-substituted tetracycline compounds include:



10 5. Methods for Synthesizing Tetracycline Compounds of the Invention


The tetracycline compounds of this invention can be synthesized using the methods described in the Schemes and/or by other techniques known to those of ordinary skill in the art.

- 15 The substituted tetracycline compounds of the invention can be synthesized using the methods described in Example 1, in the following schemes and by using art recognized techniques. All novel substituted tetracycline compounds described herein are included in the invention as compounds.



SCHEME 1


9- and 7- substituted tetracyclines can be synthesized by the method shown in Scheme 1. As shown in Scheme 1, 9- and 7-substituted tetracycline compounds can be synthesized by treating a tetracycline compound (e.g., doxycycline, 1A), with sulfuric acid and sodium nitrate. The resulting product is a mixture of the 7-nitro and 9-nitro isomers (1B and 1C, respectively). The 7-nitro (1B) and 9- nitro (1C) derivatives are treated by hydrogenation using hydrogen gas and a platinum catalyst to yield amines 1D and 1E. The isomers are separated at this time by conventional methods. To synthesize 7- or 9-substituted alkenyl derivatives, the 7- or 9-amino tetracycline compound (1E and 1F, respectively) is treated with HONO , to yield the diazonium salt (1G and 1H). The salt (1G and 1H) is treated with an appropriate reactive reagent to yield the desired compound (e.g., in Scheme 1, 7-cyclopent-1-enyl doxycycline (1H) and 9-cyclopent-1-enyl doxycycline (1I)).



SCHEME 2


As shown in Scheme 2, tetracycline compounds of the invention wherein R^7 is a carbamate or a urea derivative can be synthesized using the following protocol.

- 5 Sancycline (2A) is treated with NaNO_2 under acidic conditions forming 7-nitrosancycline (2B) in a mixture of positional isomers. 7-nitrosancycline (2B) is then treated with H_2 gas and a platinum catalyst to form the 7-amino sancycline derivative (2C). To form the urea derivative (2E), isocyanate (2D) is reacted with the 7-amino sancycline derivative (2C). To form the carbamate (2G), the appropriate acid chloride ester (2F) is reacted with 2C.



SCHEME 3


As shown in Scheme 3, tetracycline compounds of the invention, wherein R^7 is a heterocyclic (i.e. thiazole) substituted amino group can be synthesized using the above protocol. 7-amino sancycline (3A) is reacted with Fmoc-isothiocyanate (3B) to produce the protected thiourea (3C). The protected thiourea (3C) is then deprotected yielding the active sancycline thiourea (3D) compound. The sancycline thiourea (3D) is reacted with an α -haloketone (3E) to produce a thiazole substituted 7-amino sancycline (3F).



10

SCHEME 4

7- alkenyl tetracycline compounds, such as 7-alkynyl sancycline (4A) and 7-alkenyl sancycline (4B), can be hydrogenated to form 7-alkyl substituted tetracycline compounds (e.g., 7-alkyl sancycline, 4C). Scheme 4 depicts the selective hydrogenation of the 7- position double or triple bond, in saturated methanol and hydrochloric acid solution with a palladium/carbon catalyst under pressure, to yield the product.



SCHEME 5

20

In Scheme 5, a general synthetic scheme for synthesizing 7-position aryl derivatives is shown. A Suzuki coupling of an aryl boronic acid with an iodosancycline compound is shown. An iodo sancycline compound (5B) can be synthesized from sancycline by treating sancycline (5A) with at least one equivalent N-iodosuccinimide


(NIS) under acidic conditions. The reaction is quenched, and the resulting 7-iodo sancycline (5B) can then be purified using standard techniques known in the art. To form the aryl derivative, 7-iodo sancycline (5B) is treated with an aqueous base (e.g., Na_2CO_3) and an appropriate boronic acid (5C) and under an inert atmosphere. The

5 reaction is catalyzed with a palladium catalyst (e.g., $\text{Pd}(\text{OAc})_2$). The product (5D) can be purified by methods known in the art (such as HPLC). Other 7-aryl, alkenyl, and alkynyl tetracycline compounds can be synthesized using similar protocols.

The 7-substituted tetracycline compounds of the invention can also be synthesized using Stille cross couplings. Stille cross couplings can be performed using

10 an appropriate tin reagent (e.g., R-SnBu_3) and a halogenated tetracycline compound, (e.g., 7-iodosancycline). The tin reagent and the iododansancycline compound can be treated with a palladium catalyst (e.g., $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ or $\text{Pd}(\text{AsPh}_3)_2\text{Cl}_2$) and, optionally, with an additional copper salt, e.g., CuI . The resulting compound can then be purified using techniques known in the art.

15




SCHEME 6

The compounds of the invention can also be synthesized using Heck-type cross coupling reactions. As shown in Scheme 6, Heck-type cross-couplings can be

20 performed by suspending a halogenated tetracycline compound (e.g., 7-iodosancycline, 6A) and an appropriate palladium or other transition metal catalyst (e.g., $\text{Pd}(\text{OAc})_2$ and CuI) in an appropriate solvent (e.g., degassed acetonitrile). The substrate, a reactive alkene (6B) or alkyne (6D), and triethylamine are then added and the mixture is heated for several hours, before being cooled to room temperature. The resulting 7-substituted


25 alkenyl (6C) or 7-substituted alkynyl (6E) tetracycline compound can then be purified using techniques known in the art.



SCHEME 7

5

To prepare 7-(2'-Chloro-alkenyl)-tetracycline compounds, the appropriate 7-(alkynyl)-sancycline (7A) is dissolved in saturated methanol and hydrochloric acid and stirred. The solvent is then removed to yield the product (7B).




SCHEME 8

10

As depicted in Scheme 8, 5-esters of 9-substituted tetracycline compounds can be formed by dissolving the 9-substituted compounds (8A) in strong acid (e.g. HF, methanesulphonic acid, and trifluoromethanesulfonic acid) and adding the appropriate carboxylic acid to yield the corresponding esters (8B).

15

As shown in Scheme 9 below, 7 and 9 aminomethyl tetracyclines may be synthesized using reagents such as hydroxymethyl-carbamic acid benzyl ester.



SCHEME 9

The term "alkyl" includes saturated aliphatic groups, including straight-chain alkyl groups (e.g., methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, etc.), branched-chain alkyl groups (isopropyl, tert-butyl, isobutyl, etc.), cycloalkyl (alicyclic) groups (cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl), alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. The term alkyl further includes alkyl groups, which can further include oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more carbons of the hydrocarbon backbone. In certain embodiments, a straight chain or branched chain alkyl has 6 or fewer carbon atoms in its backbone (e.g., C₁-C₆ for straight chain, C₃-C₆ for branched chain), and more preferably 4 or fewer. Likewise, preferred cycloalkyls have from 3-8 carbon atoms in their ring structure, and more preferably have 5 or 6 carbons in the ring structure. The term C₁-C₆ includes alkyl groups containing 1 to 6 carbon atoms.

Moreover, the term alkyl includes both "unsubstituted alkyls" and "substituted alkyls", the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonate, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Cycloalkyls can be further substituted, e.g., with the substituents described above. An "alkylaryl" or an "arylalkyl" moiety is an alkyl substituted with an aryl (e.g., phenylmethyl (benzyl)). The term "alkyl" also includes the side chains of natural and unnatural amino acids.

The term "aryl" includes groups, including 5- and 6-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, phenyl, pyrrole, furan, thiophene, thiazole, isothiazole, imidazole, triazole, tetrazole, pyrazole, oxazole, isooxazole, pyridine, pyrazine, pyridazine, and pyrimidine, and the like. Furthermore, the term "aryl" includes multicyclic aryl groups, e.g., tricyclic, bicyclic, e.g., naphthalene, benzoxazole, benzodioxazole, benzothiazole, benzoimidazole, benzothiophene, methylenedioxyphenyl, quinoline, isoquinoline, naphthridine, indole, benzofuran, purine, benzofuran, deazapurine, or indolizine. Those

aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles", "heterocycles," "heteroaryls" or "heteroaromatics". The aromatic ring can be substituted at one or more ring positions with such substituents as described above, as for example, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, 5 alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkylaminoacarbonyl, arylalkyl aminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, arylalkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino 10 (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Aryl groups can also be fused or bridged with alicyclic or heterocyclic rings which are not aromatic so as to form a polycycle (e.g., 15 tetralin).

The term "alkenyl" includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double bond.

For example, the term "alkenyl" includes straight-chain alkenyl groups 20 (e.g., ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, etc.), branched-chain alkenyl groups, cycloalkenyl (alicyclic) groups (cyclopropenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl), alkyl or alkenyl substituted cycloalkenyl groups, and cycloalkyl or cycloalkenyl substituted alkenyl groups. The term alkenyl further includes alkenyl groups which include oxygen, 25 nitrogen, sulfur or phosphorous atoms replacing one or more carbons of the hydrocarbon backbone. In certain embodiments, a straight chain or branched chain alkenyl group has 6 or fewer carbon atoms in its backbone (e.g., C₂-C₆ for straight chain, C₃-C₆ for branched chain). Likewise, cycloalkenyl groups may have from 3-8 carbon atoms in their ring structure, and more preferably have 5 or 6 carbons in the ring structure. The 30 term C₂-C₆ includes alkenyl groups containing 2 to 6 carbon atoms.

Moreover, the term alkenyl includes both "unsubstituted alkenyls" and "substituted alkenyls", the latter of which refers to alkenyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkyl groups, alkynyl groups, halogens, hydroxyl, 35 alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl,

alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

The term "alkynyl" includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but which contain at least one triple bond.

For example, the term "alkynyl" includes straight-chain alkynyl groups (e.g., ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, etc.), branched-chain alkynyl groups, and cycloalkyl or cycloalkenyl substituted alkynyl groups. The term alkynyl further includes alkynyl groups which include oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more carbons of the hydrocarbon backbone. In certain embodiments, a straight chain or branched chain alkynyl group has 6 or fewer carbon atoms in its backbone (e.g., C₂-C₆ for straight chain, C₃-C₆ for branched chain). The term C₂-C₆ includes alkynyl groups containing 2 to 6 carbon atoms.

Moreover, the term alkynyl includes both "unsubstituted alkynyls" and "substituted alkynyls", the latter of which refers to alkynyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

Unless the number of carbons is otherwise specified, "lower alkyl" as used herein means an alkyl group, as defined above, but having from one to five carbon atoms in its backbone structure. "Lower alkenyl" and "lower alkynyl" have chain lengths of, for example, 2-5 carbon atoms.

The term "acyl" includes compounds and moieties which contain the acyl radical ($\text{CH}_3\text{CO}-$) or a carbonyl group. It includes substituted acyl moieties. The term "substituted acyl" includes acyl groups where one or more of the hydrogen atoms are replaced by for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

The term "acylamino" includes moieties wherein an acyl moiety is bonded to an amino group. For example, the term includes alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido groups.

The term "aroyl" includes compounds and moieties with an aryl or heteroaromatic moiety bound to a carbonyl group. Examples of aroyl groups include phenylcarboxy, naphthyl carboxy, etc.

The terms "alkoxyalkyl", "alkylaminoalkyl" and "thioalkoxyalkyl" include alkyl groups, as described above, which further include oxygen, nitrogen or sulfur atoms replacing one or more carbons of the hydrocarbon backbone, e.g., oxygen, nitrogen or sulfur atoms.

The term "alkoxy" includes substituted and unsubstituted alkyl, alkenyl, and alkynyl groups covalently linked to an oxygen atom. Examples of alkoxy groups include methoxy, ethoxy, isopropoxy, propoxy, butoxy, and pentoxy groups. Examples of substituted alkoxy groups include halogenated alkoxy groups. The alkoxy groups can be substituted with groups such as alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or

heteroaromatic moieties. Examples of halogen substituted alkoxy groups include, but are not limited to, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chloromethoxy, dichloromethoxy, trichloromethoxy, etc.

The term "amine" or "amino" includes compounds where a nitrogen atom is covalently bonded to at least one carbon or heteroatom. The term includes "alkyl amino" which comprises groups and compounds wherein the nitrogen is bound to at least one additional alkyl group. The term "dialkyl amino" includes groups wherein the nitrogen atom is bound to at least two additional alkyl groups. The term "arylamino" and "diarylamino" include groups wherein the nitrogen is bound to at least one or two aryl groups, respectively. The term "alkylarylamino," "alkylaminoaryl" or "arylaminoalkyl" refers to an amino group which is bound to at least one alkyl group and at least one aryl group. The term "alkaminoalkyl" refers to an alkyl, alkenyl, or alkynyl group bound to a nitrogen atom which is also bound to an alkyl group.

The term "amide," "amido" or "aminocarbonyl" includes compounds or moieties which contain a nitrogen atom which is bound to the carbon of a carbonyl or a thiocarbonyl group. The term includes "alkaminocarbonyl" or "alkylaminocarbonyl" groups which include alkyl, alkenyl, aryl or alkynyl groups bound to an amino group bound to a carbonyl group. It includes arylaminocarbonyl and arylcarbonylamino groups which include aryl or heteroaryl moieties bound to an amino group which is bound to the carbon of a carbonyl or thiocarbonyl group. The terms "alkylaminocarbonyl," "alkenylaminocarbonyl," "alkynylaminocarbonyl," "arylamino carbonyl," "alkylcarbonylamino," "alkenylcarbonylamino," "alkynylcarbonylamino," and "arylcarbonylamino" are included in term "amide." Amides also include urea groups (aminocarbonylamino) and carbamates (oxycarbonylamino).

The term "carbonyl" or "carboxy" includes compounds and moieties which contain a carbon connected with a double bond to an oxygen atom. The carbonyl can be further substituted with any moiety which allows the compounds of the invention to perform its intended function. For example, carbonyl moieties may be substituted with alkyls, alkenyls, alkynyls, aryls, alkoxy, aminos, etc. Examples of moieties which contain a carbonyl include aldehydes, ketones, carboxylic acids, amides, esters, anhydrides, etc.

The term "thiocarbonyl" or "thiocarboxy" includes compounds and moieties which contain a carbon connected with a double bond to a sulfur atom.

The term "ether" includes compounds or moieties which contain an oxygen bonded to two different carbon atoms or heteroatoms. For example, the term
5 includes "alkoxyalkyl" which refers to an alkyl, alkenyl, or alkynyl group covalently bonded to an oxygen atom which is covalently bonded to another alkyl group.

The term "ester" includes compounds and moieties which contain a carbon or a heteroatom bound to an oxygen atom which is bonded to the carbon of a carbonyl group. The term "ester" includes alkoxycarboxy groups such as
10 methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, pentoxycarbonyl, etc. The alkyl, alkenyl, or alkynyl groups are as defined above.

The term "thioether" includes compounds and moieties which contain a sulfur atom bonded to two different carbon or hetero atoms. Examples of thioethers include, but are not limited to alkthioalkyls, alkthioalkenyls, and alkthioalkynyls. The
15 term "alkthioalkyls" include compounds with an alkyl, alkenyl, or alkynyl group bonded to a sulfur atom which is bonded to an alkyl group. Similarly, the term "alkthioalkenyls" and alkthioalkynyls" refer to compounds or moieties wherein an alkyl, alkenyl, or alkynyl group is bonded to a sulfur atom which is covalently bonded to an alkynyl group.

20 The term "hydroxy" or "hydroxyl" includes groups with an -OH or -O.
The term "halogen" includes fluorine, bromine, chlorine, iodine, etc. The term "perhalogenated" generally refers to a moiety wherein all hydrogens are replaced by halogen atoms.

The terms "polycyclyl" or "polycyclic radical" refer to two or more cyclic
25 rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more carbons are common to two adjoining rings, e.g., the rings are "fused rings". Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the polycycle can be substituted with such substituents as described above, as for example, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy,
30 alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, alkylaminoacarbonyl, arylalkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, arylalkyl carbonyl, alkenylcarbonyl, aminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amido, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino
35 (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato,

sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkyl, alkylaryl, or an aromatic or heteroaromatic moiety.

The term "heteroatom" includes atoms of any element other than carbon or hydrogen. Preferred heteroatoms are nitrogen, oxygen, sulfur and phosphorus.

5 The term "prodrug moiety" includes moieties which can be metabolized *in vivo* to a hydroxyl group and moieties which may advantageously remain esterified *in vivo*. Preferably, the prodrugs moieties are metabolized *in vivo* by esterases or by other mechanisms to hydroxyl groups or other advantageous groups. Examples of prodrugs and their uses are well known in the art (See, *e.g.*, Berge et al. (1977) "Pharmaceutical
10 Salts", *J. Pharm. Sci.* 66:1-19). The prodrugs can be prepared *in situ* during the final isolation and purification of the compounds, or by separately reacting the purified compound in its free acid form or hydroxyl with a suitable esterifying agent. Hydroxyl groups can be converted into esters *via* treatment with a carboxylic acid. Examples of prodrug moieties include substituted and unsubstituted, branch or unbranched lower
15 alkyl ester moieties, (*e.g.*, propionic acid esters), lower alkenyl esters, di-lower alkyl-amino lower-alkyl esters (*e.g.*, dimethylaminoethyl ester), acylamino lower alkyl esters (*e.g.*, acetyloxymethyl ester), acyloxy lower alkyl esters (*e.g.*, pivaloyloxymethyl ester), aryl esters (phenyl ester), aryl-lower alkyl esters (*e.g.*, benzyl ester), substituted (*e.g.*, with methyl, halo, or methoxy substituents) aryl and aryl-lower alkyl esters, amides,
20 lower-alkyl amides, di-lower alkyl amides, and hydroxy amides. Preferred prodrug moieties are propionic acid esters and acyl esters.

It will be noted that the structure of some of the tetracycline compounds of this invention includes asymmetric carbon atoms. It is to be understood accordingly that the isomers arising from such asymmetry (*e.g.*, all enantiomers and diastereomers)
25 are included within the scope of this invention, unless indicated otherwise. Such isomers can be obtained in substantially pure form by classical separation techniques and by stereochemically controlled synthesis. Furthermore, the structures and other compounds and moieties discussed in this application also include all tautomers thereof.

30 6. Methods for Treating Tetracycline Responsive States

The invention also pertains to methods for treating a tetracycline responsive states in subjects, by administering to a subject an effective amount of a tetracycline compound of the invention (*e.g.*, a compound of Formula I, II, III, IV, or otherwise described herein), such that the tetracycline responsive state is treated.

The term "treating" includes curing as well as ameliorating at least one symptom of the state, disease or disorder, e.g., the tetracycline compound responsive state.

- The language "tetracycline compound responsive state" or "tetracycline responsive state" includes states which can be treated, prevented, or otherwise ameliorated by the administration of a tetracycline compound of the invention, e.g., a 3, 10, and/or 12a substituted tetracycline compound. Tetracycline compound responsive states include bacterial, viral, and fungal infections (including those which are resistant to other tetracycline compounds), cancer (e.g., prostate, breast, colon, lung melanoma and lymph cancers and other disorders characterized by unwanted cellular proliferation, including, but not limited to, those described in U.S. 6,100,248), arthritis, osteoporosis, diabetes, and other states for which tetracycline compounds have been found to be active (see, for example, U.S. Patent Nos. 5,789,395; 5,834,450; 6,277,061 and 5,532,227, each of which is expressly incorporated herein by reference).
- Compounds of the invention can be used to prevent or control important mammalian and veterinary diseases such as diarrhea, urinary tract infections, infections of skin and skin structure, ear, nose and throat infections, wound infection, mastitis and the like. In addition, methods for treating neoplasms using tetracycline compounds of the invention are also included (van der Bozert *et al.*, *Cancer Res.*, 48:6686-6690 (1988)). In a further embodiment, the tetracycline responsive state is not a bacterial infection. In another embodiment, the tetracycline compounds of the invention are essentially non-antibacterial. For example, non-antibacterial tetracycline compounds of the invention may have MIC values greater than about 4 µg/ml (as measured by assays known in the art and/or the assay given in Example 2).
- Tetracycline compound responsive states also include inflammatory process associated states (IPAS). The term "inflammatory process associated state" includes states in which inflammation or inflammatory factors (e.g., matrix metalloproteinases (MMPs), nitric oxide (NO), TNF, interleukins, plasma proteins, cellular defense systems, cytokines, lipid metabolites, proteases, toxic radicals, adhesion molecules, *etc.*) are involved or are present in an area in aberrant amounts, e.g., in amounts which may be advantageous to alter, e.g., to benefit the subject. The inflammatory process is the response of living tissue to damage. The cause of inflammation may be due to physical damage, chemical substances, micro-organisms, tissue necrosis, cancer or other agents. Acute inflammation is short-lasting, lasting only a few days. If it is longer lasting however, then it may be referred to as chronic inflammation.

IPAF's include inflammatory disorders. Inflammatory disorders are generally characterized by heat, redness, swelling, pain and loss of function. Examples of causes of inflammatory disorders include, but are not limited to, microbial infections (e.g., bacterial and fungal infections), physical agents (e.g., burns, radiation, and trauma),
5 chemical agents (e.g., toxins and caustic substances), tissue necrosis and various types of immunologic reactions.

Examples of inflammatory disorders include, but are not limited to, osteoarthritis, rheumatoid arthritis, acute and chronic infections (bacterial and fungal, including diphtheria and pertussis); acute and chronic bronchitis, sinusitis, and upper
10 respiratory infections, including the common cold; acute and chronic gastroenteritis and colitis; acute and chronic cystitis and urethritis; acute and chronic dermatitis; acute and chronic conjunctivitis; acute and chronic serositis (pericarditis, peritonitis, synovitis, pleuritis and tendinitis); uremic pericarditis; acute and chronic cholecystitis; acute and chronic vaginitis; acute and chronic uveitis; drug reactions; insect bites; burns (thermal,
15 chemical, and electrical); and sunburn.

Tetracycline compound responsive states also include NO associated states. The term "NO associated state" includes states which involve or are associated with nitric oxide (NO) or inducible nitric oxide synthase (iNOS). NO associated state includes states which are characterized by aberrant amounts of NO and/or iNOS. Preferably, the
20 NO associated state can be treated by administering tetracycline compounds of the invention, e.g., a 3, 10, and/or 12a substituted tetracycline compound. The disorders, diseases and states described in U.S. Patents Nos. 6,231,894; 6,015,804; 5,919,774; and 5,789,395 are also included as NO associated states. The entire contents of each of these patents are hereby incorporated herein by reference.

25 Other examples of NO associated states include, but are not limited to, malaria, senescence, diabetes, vascular stroke, neurodegenerative disorders (Alzheimer's disease & Huntington's disease), cardiac disease (reperfusion-associated injury following infarction), juvenile diabetes, inflammatory disorders, osteoarthritis, rheumatoid arthritis, acute, recurrent and chronic infections (bacterial, viral and fungal); acute and
30 chronic bronchitis, sinusitis, and respiratory infections, including the common cold; acute and chronic gastroenteritis and colitis; acute and chronic cystitis and urethritis; acute and chronic dermatitis; acute and chronic conjunctivitis; acute and chronic serositis (pericarditis, peritonitis, synovitis, pleuritis and tendonitis); uremic pericarditis; acute and chronic cholecystitis; cystic fibrosis, acute and chronic vaginitis; acute and
35 chronic uveitis; drug reactions; insect bites; burns (thermal, chemical, and electrical); and sunburn.

The term "inflammatory process associated state" also includes, in one embodiment, matrix metalloproteinase associated states (MMPAS). MMPAS include states characterized by aberrant amounts of MMPs or MMP activity. These are also include as tetracycline compound responsive states which may be treated using
5 compounds of the invention, e.g., 3, 10, and/or 12a substituted tetracycline compounds.

Examples of matrix metalloproteinase associated states ("MMPAS's") include, but are not limited to, arteriosclerosis, corneal ulceration, emphysema, osteoarthritis, multiple sclerosis(Liedtke *et al.*, *Ann. Neurol.* 1998, 44:35-46; Chandler *et al.*, *J. Neuroimmunol.* 1997, 72:155-71), osteosarcoma, osteomyelitis, bronchiectasis, chronic
10 pulmonary obstructive disease, skin and eye diseases, periodontitis, osteoporosis, rheumatoid arthritis, ulcerative colitis, inflammatory disorders, tumor growth and invasion (Stetler-Stevenson *et al.*, *Annu. Rev. Cell Biol.* 1993, 9:541-73; Tryggvason *et al.*, *Biochim. Biophys. Acta* 1987, 907:191-217; Li *et al.*, *Mol. Carcinog.* 1998, 22:84-89)),metastasis, acute lung injury, stroke, ischemia, diabetes, aortic or vascular
15 aneurysms, skin tissue wounds, dry eye, bone and cartilage degradation (Greenwald *et al.*, *Bone* 1998, 22:33-38; Ryan *et al.*, *Curr. Op. Rheumatol.* 1996, 8:238-247). Other MMPAS include those described in U.S. Pat. Nos. 5,459,135; 5,321,017; 5,308,839; 5,258,371; 4,935,412; 4,704,383, 4,666,897, and RE 34,656, incorporated herein by reference in their entirety.

20 In another embodiment, the tetracycline compound responsive state is cancer. Examples of cancers which the tetracycline compounds of the invention may be useful to treat include all solid tumors, i.e., carcinomas e.g., adenocarcinomas, and sarcomas. Adenocarcinomas are carcinomas derived from glandular tissue or in which the tumor cells form recognizable glandular structures. Sarcomas broadly include tumors whose
25 cells are embedded in a fibrillar or homogeneous substance like embryonic connective tissue. Examples of carcinomas which may be treated using the methods of the invention include, but are not limited to, carcinomas of the prostate, breast, ovary, testis, lung, colon, and breast. The methods of the invention are not limited to the treatment of these tumor types, but extend to any solid tumor derived from any organ system.
30 Examples of treatable cancers include, but are not limited to, colon cancer, bladder cancer, breast cancer, melanoma, ovarian carcinoma, prostatic carcinoma, lung cancer, and a variety of other cancers as well. The methods of the invention also cause the inhibition of cancer growth in adenocarcinomas, such as, for example, those of the prostate, breast, kidney, ovary, testes, and colon.

35

In an embodiment, the tetracycline responsive state of the invention is cancer. The invention pertains to a method for treating a subject suffering or at risk of suffering from cancer, by administering an effective amount of a substituted tetracycline compound, such that inhibition cancer cell growth occurs, i.e., cellular proliferation, 5 invasiveness, metastasis, or tumor incidence is decreased, slowed, or stopped. The inhibition may result from inhibition of an inflammatory process, down-regulation of an inflammatory process, some other mechanism, or a combination of mechanisms. Alternatively, the tetracycline compounds may be useful for preventing cancer recurrence, for example, to treat residual cancer following surgical resection or radiation 10 therapy. The tetracycline compounds useful according to the invention are especially advantageous as they are substantially non-toxic compared to other cancer treatments. In a further embodiment, the compounds of the invention are administered in combination with standard cancer therapy, such as, but not limited to, chemotherapy.

Examples of tetracycline responsive states also include neurological disorders 15 which include both neuropsychiatric and neurodegenerative disorders, but are not limited to, such as Alzheimer's disease, dementias related to Alzheimer's disease (such as Pick's disease), Parkinson's and other Lewy diffuse body diseases, senile dementia, Huntington's disease, Gilles de la Tourette's syndrome, multiple sclerosis, amyotrophic lateral sclerosis (ALS), progressive supranuclear palsy, epilepsy, and Creutzfeldt-Jakob 20 disease; autonomic function disorders such as hypertension and sleep disorders, and neuropsychiatric disorders, such as depression, schizophrenia, schizoaffective disorder, Korsakoff's psychosis, mania, anxiety disorders, or phobic disorders; learning or memory disorders, *e.g.*, amnesia or age-related memory loss, attention deficit disorder, dysthymic disorder, major depressive disorder, mania, obsessive-compulsive disorder, 25 psychoactive substance use disorders, anxiety, phobias, panic disorder, as well as bipolar affective disorder, *e.g.*, severe bipolar affective (mood) disorder (BP-1), bipolar affective neurological disorders, *e.g.*, migraine and obesity. Further neurological disorders include, for example, those listed in the American Psychiatric Association's Diagnostic and Statistical manual of Mental Disorders (DSM), the most current version 30 of which is incorporated herein by reference in its entirety.

The language "in combination with" another therapeutic agent or treatment includes co-administration of the tetracycline compound, (*e.g.*, inhibitor) and with the other therapeutic agent or treatment, administration of the tetracycline compound first, 35 followed by the other therapeutic agent or treatment and administration of the other therapeutic agent or treatment first, followed by the tetracycline compound. The other therapeutic agent may be any agent which is known in the art to treat, prevent, or reduce

the symptoms of an IPAS. Furthermore, the other therapeutic agent may be any agent of benefit to the patient when administered in combination with the administration of an tetracycline compound. In one embodiment, the cancers treated by methods of the invention include those described in U.S. Patent Nos. 6,100,248; 5,843,925; 5,837,696; 5 or 5,668,122, incorporated herein by reference in their entirety.

In another embodiment, the tetracycline compound responsive state is diabetes, e.g., juvenile diabetes, diabetes mellitus, diabetes type I, or diabetes type II. In a further embodiment, protein glycosylation is not affected by the administration of the tetracycline compounds of the invention. In another embodiment, the tetracycline 10 compound of the invention is administered in combination with standard diabetic therapies, such as, but not limited to insulin therapy. In a further embodiment, the IPAS includes disorders described in U.S. Patents Nos. 5,929,055; and 5,532,227, incorporated herein by reference in their entirety.

In another embodiment, the tetracycline compound responsive state is a bone 15 mass disorder. Bone mass disorders include disorders where a subject's bones are disorders and states where the formation, repair or remodeling of bone is advantageous. For examples bone mass disorders include osteoporosis (e.g., a decrease in bone strength and density), bone fractures, bone formation associated with surgical procedures (e.g., facial reconstruction), osteogenesis imperfecta (brittle bone disease), hypophosphatasia, 20 Paget's disease, fibrous dysplasia, osteopetrosis, myeloma bone disease, and the depletion of calcium in bone, such as that which is related to primary hyperparathyroidism. Bone mass disorders include all states in which the formation, repair or remodeling of bone is advantageous to the subject as well as all other disorders associated with the bones or skeletal system of a subject which can be treated with the 25 tetracycline compounds of the invention. In a further embodiment, the bone mass disorders include those described in U.S. Patents Nos. 5,459,135; 5,231,017; 5,998,390; 5,770,588; RE 34,656; 5,308,839; 4,925,833; 3,304,227; and 4,666,897, each of which is hereby incorporated herein by reference in its entirety.

In another embodiment, the tetracycline compound responsive state is acute lung 30 injury. Acute lung injuries include adult respiratory distress syndrome (ARDS), post-pump syndrome (PPS), and trauma. Trauma includes any injury to living tissue caused by an extrinsic agent or event. Examples of trauma include, but are not limited to, crush injuries, contact with a hard surface, or cutting or other damage to the lungs.

35

The invention also pertains to a method for treating acute lung injury by administering a substituted tetracycline compound of the invention.

The tetracycline responsive states of the invention also include chronic lung disorders. The invention pertains to methods for treating chronic lung disorders by administering a tetracycline compound, such as those described herein. The method includes administering to a subject an effective amount of a substituted tetracycline compound such that the chronic lung disorder is treated. Examples of chronic lung disorders include, but are not limited, to asthma, cystic fibrosis, and emphysema. In a further embodiment, the tetracycline compounds of the invention used to treat acute and/or chronic lung disorders such as those described in U.S. Patents No. 5,977,091; 6,043,231; 5,523,297; and 5,773,430, each of which is hereby incorporated herein by reference in its entirety.

In yet another embodiment, the tetracycline compound responsive state is ischemia, stroke, or ischemic stroke. The invention also pertains to a method for treating ischemia, stroke, or ischemic stroke by administering an effective amount of a substituted tetracycline compound of the invention. In a further embodiment, the tetracycline compounds of the invention are used to treat such disorders as described in U.S. Patents No. 6,231,894; 5,773,430; 5,919,775 or 5,789,395, incorporated herein by reference.

In another embodiment, the tetracycline compound responsive state is a skin wound. The invention also pertains, at least in part, to a method for improving the healing response of the epithelialized tissue (e.g., skin, mucosae) to acute traumatic injury (e.g., cut, burn, scrape, etc.). The method may include using a tetracycline compound of the invention (which may or may not have antibacterial activity) to improve the capacity of the epithelialized tissue to heal acute wounds. The method may increase the rate of collagen accumulation of the healing tissue. The method may also decrease the proteolytic activity in the epithelialized tissue by decreasing the collagenolytic and/or gelatinolytic activity of MMPs. In a further embodiment, the tetracycline compound of the invention is administered to the surface of the skin (e.g., topically). In a further embodiment, the tetracycline compound of the invention used to treat a skin wound, and other such disorders as described in, for example, U.S. Patent Nos. 5,827,840; 4,704,383; 4,935,412; 5,258,371; 5,308,839; 5,459,135; 5,532,227; and 6,015,804; each of which is incorporated herein by reference in its entirety.

In yet another embodiment, the tetracycline compound responsive state is an aortic or vascular aneurysm in vascular tissue of a subject (e.g., a subject having or at risk of having an aortic or vascular aneurysm, etc.). The tetracycline compound may by

effective to reduce the size of the vascular aneurysm or it may be administered to the subject prior to the onset of the vascular aneurysm such that the aneurysm is prevented. In one embodiment, the vascular tissue is an artery, e.g., the aorta, e.g., the abdominal aorta. In a further embodiment, the tetracycline compounds of the invention are used to
5 treat disorders described in U.S. Patent Nos. 6,043,225 and 5,834,449, incorporated herein by reference in their entirety.

Bacterial infections may be caused by a wide variety of gram positive and gram negative bacteria. The compounds of the invention are useful as antibiotics against organisms which are resistant to other tetracycline compounds. The antibiotic activity
10 of the tetracycline compounds of the invention may be determined using the method discussed in Example 2, or by using the in vitro standard broth dilution method described in Waitz, J.A., *National Commission for Clinical Laboratory Standards, Document M7-A2*, vol. 10, no. 8, pp. 13-20, 2nd edition, Villanova, PA (1990).

The tetracycline compounds may also be used to treat infections traditionally
15 treated with tetracycline compounds such as, for example, rickettsiae; a number of gram-positive and gram-negative bacteria; and the agents responsible for lymphogranuloma venereum, inclusion conjunctivitis, psittacosis. The tetracycline compounds may be used to treat infections of, e.g., *K. pneumoniae*, *Salmonella*, *E. hirae*, *A. baumannii*, *B. catarrhalis*, *H. influenzae*, *P. aeruginosa*, *E. faecium*, *E. coli*, *S.*
20 *aureus* or *E. faecalis*. In one embodiment, the tetracycline compound is used to treat a bacterial infection that is resistant to other tetracycline antibiotic compounds. The tetracycline compound of the invention may be administered with a pharmaceutically acceptable carrier.

The language "effective amount" of the compound is that amount necessary or
25 sufficient to treat or prevent a tetracycline compound responsive state. The effective amount can vary depending on such factors as the size and weight of the subject, the type of illness, or the particular tetracycline compound. For example, the choice of the tetracycline compound can affect what constitutes an "effective amount". One of ordinary skill in the art would be able to study the aforementioned factors and make the
30 determination regarding the effective amount of the tetracycline compound without undue experimentation.

The invention also pertains to methods of treatment against microorganism infections and associated diseases. The methods include administration of an effective amount of one or more tetracycline compounds to a subject. The subject can be either a
35 plant or, advantageously, an animal, e.g., a mammal, e.g., a human.

In the therapeutic methods of the invention, one or more tetracycline compounds of the invention may be administered alone to a subject, or more typically a compound of the invention will be administered as part of a pharmaceutical composition in mixture with conventional excipient, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, oral or other desired administration and which do not deleteriously react with the active compounds and are not deleterious to the recipient thereof.

7. **Pharmaceutical Compositions of the Invention**

The invention also pertains to pharmaceutical compositions comprising a therapeutically effective amount of a tetracycline compound (e.g., a compound of Formula (I), (II), or (III), or any other compound described herein) and, optionally, a pharmaceutically acceptable carrier.

The language "pharmaceutically acceptable carrier" includes substances capable of being coadministered with the tetracycline compound(s), and which allow both to perform their intended function, e.g., treat or prevent a tetracycline responsive state. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with the active compounds of the invention.

The tetracycline compounds of the invention that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of the tetracycline compounds of the invention that are basic in nature are those that form non-toxic acid addition salts, i.e., salts containing pharmaceutically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, acid citrate, tartrate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucaronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and palmoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts. Although such salts must be

pharmaceutically acceptable for administration to a subject, e.g., a mammal, it is often desirable in practice to initially isolate a tetracycline compound of the invention from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and
5 subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the
10 desired solid salt is readily obtained. The preparation of other tetracycline compounds of the invention not specifically described in the foregoing experimental section can be accomplished using combinations of the reactions described above that will be apparent to those skilled in the art.

The preparation of other tetracycline compounds of the invention not
15 specifically described in the foregoing experimental section can be accomplished using combinations of the reactions described above that will be apparent to those skilled in the art.

The tetracycline compounds of the invention that are acidic in nature are capable of forming a wide variety of base salts. The chemical bases that may be used as
20 reagents to prepare pharmaceutically acceptable base salts of those tetracycline compounds of the invention that are acidic in nature are those that form non-toxic base salts with such compounds. Such non-toxic base salts include, but are not limited to those derived from such pharmaceutically acceptable cations such as alkali metal cations (e.g., potassium and sodium) and alkaline earth metal cations (e.g., calcium and
25 magnesium), ammonium or water-soluble amine addition salts such as N-methylglucamine (meglumine), and the lower alkanolammonium and other base salts of pharmaceutically acceptable organic amines. The pharmaceutically acceptable base addition salts of tetracycline compounds of the invention that are acidic in nature may be formed with pharmaceutically acceptable cations by conventional methods. Thus, these
30 salts may be readily prepared by treating the tetracycline compound of the invention with an aqueous solution of the desired pharmaceutically acceptable cation and evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, a lower alkyl alcohol solution of the tetracycline compound of the invention may be mixed with an alkoxide of the desired metal and the solution
35 subsequently evaporated to dryness.

The preparation of other tetracycline compounds of the invention not specifically described in the foregoing experimental section can be accomplished using combinations of the reactions described above that will be apparent to those skilled in the art.

- 5 The tetracycline compounds of the invention and pharmaceutically acceptable salts thereof can be administered via either the oral, parenteral or topical routes. In general, these compounds are most desirably administered in effective dosages, depending upon the weight and condition of the subject being treated and the particular route of administration chosen. Variations may occur depending upon the
- 10 species of the subject being treated and its individual response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval at which such administration is carried out.

- The pharmaceutical compositions of the invention may be administered alone or in combination with other known compositions for treating tetracycline
- 15 responsive states in a subject, e.g., a mammal. Preferred mammals include pets (e.g., cats, dogs, ferrets, etc.), farm animals (cows, sheep, pigs, horses, goats, etc.), lab animals (rats, mice, monkeys, etc.), and primates (chimpanzees, humans, gorillas). The language "in combination with" a known composition is intended to include simultaneous administration of the composition of the invention and the known composition,
- 20 administration of the composition of the invention first, followed by the known composition and administration of the known composition first, followed by the composition of the invention. Any of the therapeutically composition known in the art for treating tetracycline responsive states can be used in the methods of the invention.

- The tetracycline compounds of the invention may be administered alone
- 25 or in combination with pharmaceutically acceptable carriers or diluents by any of the routes previously mentioned, and the administration may be carried out in single or multiple doses. For example, the novel therapeutic agents of this invention can be administered advantageously in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of
- 30 tablets, capsules, lozenges, troches, hard candies, powders, sprays (e.g., aerosols, etc.), creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored.
- 35 In general, the therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with
5 granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols.
10 When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof. The compositions of the invention may be formulated such that
15 the tetracycline compositions are released over a period of time after administration.

For parenteral administration (including intraperitoneal, subcutaneous, intravenous, intradermal or intramuscular injection), solutions of a therapeutic compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered
20 (preferably pH greater than 8) if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the
25 art. For parenteral application, examples of suitable preparations include solutions, preferably oily or aqueous solutions as well as suspensions, emulsions, or implants, including suppositories. Therapeutic compounds may be formulated in sterile form in multiple or single dose formats such as being dispersed in a fluid carrier such as sterile physiological saline or 5% saline dextrose solutions commonly used with injectables.

30 Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin. Examples of methods of topical administration include transdermal, buccal or sublingual application. For topical applications, therapeutic compounds can be suitably admixed in a pharmacologically inert topical carrier such as a gel, an ointment, a lotion or a cream.
35 Such topical carriers include water, glycerol, alcohol, propylene glycol, fatty alcohols, triglycerides, fatty acid esters, or mineral oils. Other possible topical carriers are liquid

petrolatum, isopropylpalmitate, polyethylene glycol, ethanol 95%, polyoxyethylene monolaurate 5% in water, sodium lauryl sulfate 5% in water, and the like. In addition, materials such as anti-oxidants, humectants, viscosity stabilizers and the like also may be added if desired.

5 For enteral application, particularly suitable are tablets, dragees or capsules having talc and/or carbohydrate carrier binder or the like, the carrier preferably being lactose and/or corn starch and/or potato starch. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including those wherein the active component is protected with differentially
10 degradable coatings, e.g., by microencapsulation, multiple coatings, etc.

In addition to treatment of human subjects, the therapeutic methods of the invention also will have significant veterinary applications, e.g. for treatment of livestock such as cattle, sheep, goats, cows, swine and the like; poultry such as chickens, ducks, geese, turkeys and the like; horses; and pets such as dogs and cats. Also, the
15 compounds of the invention may be used to treat non-animal subjects, such as plants.

It will be appreciated that the actual preferred amounts of active compounds used in a given therapy will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application, the particular site of administration, etc. Optimal administration rates for a given protocol of
20 administration can be readily ascertained by those skilled in the art using conventional dosage determination tests conducted with regard to the foregoing guidelines.

In general, compounds of the invention for treatment can be administered to a subject in dosages used in prior tetracycline therapies. See, for example, the *Physicians' Desk Reference*. For example, a suitable effective dose of one or more
25 compounds of the invention will be in the range of from 0.01 to 100 milligrams per kilogram of body weight of recipient per day, preferably in the range of from 0.1 to 50 milligrams per kilogram body weight of recipient per day, more preferably in the range of 1 to 20 milligrams per kilogram body weight of recipient per day. The desired dose is suitably administered once daily, or several sub-doses, e.g. 2 to 5 sub-doses, are
30 administered at appropriate intervals through the day, or other appropriate schedule. It will also be understood that normal, conventionally known precautions will be taken regarding the administration of tetracyclines generally to ensure their efficacy under normal use circumstances. Especially when employed for therapeutic treatment of humans and animals *in vivo*, the practitioner should take all sensible precautions to
35 avoid conventionally known contradictions and toxic effects. Thus, the conventionally recognized adverse reactions of gastrointestinal distress and inflammations, the renal

toxicity, hypersensitivity reactions, changes in blood, and impairment of absorption through aluminum, calcium, and magnesium ions should be duly considered in the conventional manner.

Furthermore, the invention also pertains to the use of a tetracycline compound of formula I, II, III, IV, or any other compound described herein, for the preparation of a medicament. The medicament may include a pharmaceutically acceptable carrier and the tetracycline compound is an effective amount, e.g., an effective amount to treat a tetracycline responsive state.

10 EXEMPLIFICATION OF THE INVENTION

Compounds of the invention may be made as described below, with modifications to the procedure below within the skill of those of ordinary skill in the art.

EXAMPLE 1: Synthesis of Substituted Tetracycline Compounds

15

7 Iodo Sancycline

One gram of sancycline was dissolved in 25 mL of TFA (trifluoroacetic acid) that was cooled to 0 C (on ice). 1.2 equivalents of N-iodosuccinimide (NIS) was added to the reaction mixture and reacted for forty minutes. The reaction was removed from the ice bath and was allowed to react at room temperature for an additional five hours. The mixture was then analyzed by HPLC and TLC, was driven to completion by the stepwise addition of NIS. After completion of the reaction, the TFA was removed *in vacuo* and 3 mL of MeOH was added to dissolve the residue. The methanolic solution was added slowly to a rapidly stirring solution of diethyl ether to form a greenish brown precipitate. The 7-iodo isomer of sancycline was purified by treating the 7-iodo product with activated charcoal, filtering through Celite, and subsequent removal of the solvent *in vacuo* to produce the 7-isomer compound as a pure yellow solid in 75% yield. MS(M+H) (formic acid solvent) 541.3.

25
Rt: Hypersil C18 BDS Column, 11.73

30 ¹H NMR (Methanol d₄-300 MHz) δ 7.87-7.90 (d, 1H), 6.66-6.69 (d, 1H), 4.06 (s, 1H), 2.98 (s, 6H), 2.42 (m, 1H), 2.19 (m, 1H), 1.62 (m, 4H), 0.99 (m, 2H)

7-Phenyl Sancycline

7-iodosancycline, 150 mg (0.28 mM), Pd(OAc)₂ and 10 mL of MeOH are added to a flask with a stir bar and the system degassed 3x using argon. Na₂CO₃ (87 mg, 0.8 mM) dissolved in water and argon degassed is added via syringe is added along

with phenylboronic acid (68 mg, 0.55 mM) in MeOH that was also degassed. The reaction was followed by HPLC for 2 hours and cooled to room temperature. The solution was filtered, and dried to produce a crude mixture. The solid was dissolved in dimethylformamide and injected onto a preparative HPLC system using C18 reverse-phase silica. The fraction at 36-38 minutes was isolated, and the solvent removed in vacuo to yield the product plus salts. The salts were removed by extraction into 50:25:25 water, butanol, ethyl acetate and dried *in vacuo*. This solid was dissolved in MeOH and the HCl salt made by bubbling in HCl gas. The solvent was removed to produce the product in 42% yield as a yellow solid.

10

Rt 21.6 min: MS (M+H, formic acid solvent): 491.3

¹H NMR (Methanol d₄-300 MHz) δ 7.87 (d, J=8.86 Hz, 1H), 7.38 (m, 5H), 6.64 (d, 8.87 Hz, 1H), 4.00 (s, 1H), 3.84 (s, 2H), 3.01 (s, 6H), 2.46 (m, 2H), 1.63 (m, 4H), 0.95 (m, 2H)

15

7-(4'-Iodo-1',3'-carboethoxy-1',3'-butadiene) Sancycline

7-I-Sancycline (1 gm, 1.86 mmol), was dissolved in 25 mL of acetonitrile and was degassed and purged with nitrogen (three times). To this suspension Pd(OAc)₂ (20 mg, .089 mmol), CuI (10 mg, .053 mmol), (o-tolyl)₃P (56 mg, .183 mmol) were added and purged with nitrogen. Ethyl propiolate (1 mL) and triethylamine (1 mL) were added to the suspension. It turned to a brown solution upon addition of Et₃N. The reaction mixture was then heated to 70 degrees C for two hours. Progress of the reaction was monitored by HPLC. It was then cooled down to room temperature and was filtered through celite. Evaporation of the solvent gave a brown solid, which was then purified on preparative HPLC to give a yellow solid.

25

7-(2'-Chloroethenyl)-Sancycline

To a solution/suspension of 0.65 g (1 mmol) of 7-iodo sancycline, 0.05 g tetrakis triphenyl phosphinato palladate, 0.012 g palladium acetate, 0.05 g copper (I) iodide in 10 mL acetonitrile, 2 mL triethylamine and 0.5 g trimethylsilyl acetylene was added at room temperature. The reaction proceeded for two hours before being filtered through a celite bed and concentrated. The crude product was purified by preparative HPLC. The collected fractions were concentrated and the residue was taken up in about 1 mL of methanol and 2 mL of HCl saturated methanol. The product was precipitated with ether. The solids were filtered off and dried under reduced pressure. NMR


35

spectroscopy and LC-MS showed that the compound was 7-(2-chloroethyl) sancycline.

7-(4'-Aminophenyl) Sancycline

- 5 To a solution of 200 mg of 7-(4-nitrophenyl) sancycline in 50 mL methanol, 10 mg of 10% palladium on charcoal catalyst was added. The reaction mixture was shaken under 40 psi hydrogen pressure for 2 hours and was then filtered followed by concentration. The residue was further purified by preparative HPLC. 35 mg was isolated as the HCl salt and the structure was proved by MNR and LC-MS to be
- 10 7-(4-aminophenyl) sancycline.


7-(NN-Dimethylpropynyl)-Sancycline




- 7-I-Sancycline (1 gm, 1.86 mmol), taken in 25 mL of acetonitrile was
- 15 degassed and purged with nitrogen (three times). To this suspension $\text{Pd}(\text{OAc})_2$ (20 mg, .089 mmol), CuI (10 mg, .053 mmol), $(o\text{-tolyl})_3\text{P}$ (56 mg, 0.183 mmol) were added and purged with nitrogen for few minutes. NN-Dimethylpropyne (308 mg, 3.72 mmol) and triethylamine (1 mL) were added to the suspension. It was turned into a brown solution upon addition of Et_3N . The reaction mixture was then heated to 70 °C for 3 hours.
- 20 Progress of the reaction was monitored by HPLC. It was then cooled down to room temperature and was filtered through celite. Evaporation of the solvent gave a brown solid, which was then purified on preparative HPLC to give a yellow solid. The structure of this compound has been characterized using ^1H NMR, HPLC, and MS.

25

30

7-(2'-Chloro-3-Hydroxypropenyl)-Sancycline

7-(alkynyl)-sancycline (100 mg) was taken in 20 ml of saturated MeOH/HCl and
 5 stirred for 20 min. The solvent was then evaporated to give a yellow powder. The
 structure of this compound has been characterized using ¹H NMR, HPLC, and MS.


7-(3'-Methoxyphenylethyl)-Sancycline

10

7-(3'-Methoxyphenylethynyl)-sancycline (1mmol)/ was taken in saturated
 solution of MeOH/HCl. To this solution 10% Pd/C was added and was subjected to
 hydrogenation at 50 psi for 12 hrs. It was then filtered through celite. The solvent was
 15 evaporated to give a yellow powder. Finally, it was precipitated from
 MeOH/diethylether. The structure of this compound has been characterized using ¹H
 NMR, HPLC, and MS.

(2-Dimethylamino-Acetyl-amino)-Sancycline


20




NN-Dimethylglycine (1.2 mmol) was dissolved in DMF (5 mL) and *O*-
 Benzotriazol-1-yl-*N, N, N', N'*-tetramethyluronium hexafluorophosphate (HBTU, 1.2
 mmol) was added. The solution was then stirred for 5 minutes at room temperature. To
 25 this solution, 7-aminosancycline (1 mmol) was added, followed by the addition of

diisopropylethyl amine (DIEA, 1.2 mmol). The reaction was then stirred at room temperature for 2 hours. The solvent, DMF, was removed on vacuum. The crude material was dissolved in 5 mL of MeOH and filtered using autovials and purified using preparative HPLC. The structure of the product has been characterized using ¹H NMR, HPLC, and MS.

7-(N-Methylsulphonamidopropargylamine) Sancycline




To a mixture of 7-iodosancycline mono trifluoroacetic acid salt (1g; 1.53 mmols), palladium II acetate (17.2 mg; 0.076 mmols), tetrakis triphenylphosphine palladium (176.8 mg; 0.153 mmols), and copper (I) iodide (49 mg; 0.228 mmols) was added 15 ml of reagent grade acetonitrile in a clean dry 2 necked round bottom flask. The reaction was purged with a slow stream of argon gas, with stirring, for 5 minutes before the addition (in one portion as a solid) of N-methylsulphonamidopropargyl amine. The sulphonamide was prepared by a method known in the art (J.Med.Chem 31(3) 1988; 577-82). This was followed by one milliliter of triethylamine (1 ml; 0.726 mmols) and the reaction was stirred, under an argon atmosphere, for approximately 1.0 hour at ambient temperature. The reaction mixture was suctioned filtered through a pad of diatomaceous earth and washed with acetonitrile. The filtrates were reduced to dryness under vacuo and the residue was treated with a dilute solution of trifluoroacetic acid in acetonitrile to adjust the pH to approximately 2. The residue was treated with more dilute trifluoroacetic acid in acetonitrile, resulting in the formation of a precipitate, which was removed via suction filtration. The crude filtrates were purified utilizing reverse phase HPLC with DVB as the solid phase; and a gradient of 1:1 methanol/acetonitrile 1% trifluoroacetic acid and 1% trifluoroacetic acid in water. The appropriate fractions were reduced to dryness under reduced pressure and solid collected. The product was characterized via ¹H NMR, mass spectrogram and LC reverse phase.

7-(2'-methoxy-5'-formylphenyl)sancycline

7-iodo-sancycline (1g, 1.53 mmol), Pd(OAc)₂ (34 mg, 0.153 mmol), and MeOH (50 mL) were combined in a 250 mL 2 neck round bottom flask equipped with a condenser and argon line. The solution was then purged with argon (15min) while heated in an oil bath to approximately 70°C. Sodium carbonate (482mg, 4.58mmol) was dissolved in water (3-5mL) and added to reaction flask. The flask was then purged with argon for another 5 minutes. 2-Methoxy-5-formylphenyl boronic acid (333mg, 1.83mmol) was dissolved in MeOH (5mL) and added to reaction flask. The flask was then purged again with argon for 10 minutes. The reaction was monitored to completion within 3 hours. The contents of the flask were filtered through filter paper and the remaining solvent was evacuated. To make the hydrochloric acid salt, the residue was dissolved in MeOH (sat. HCl) to make the HCl salt. The solution was then filtered and the solvent was evacuated. The product was then characterized by ¹H NMR, LC-MS.


15

7-(2'-Methoxy-5'-N,N'-Dimethylaminomethylphenyl)Sancycline

7-(2'-methoxy-5'-formylphenyl)sancycline (1g, 1.82mmol), dimethylamine HCl (297 mg, 3.64 mmol), triethylamine (506 μ L, 3.64 mmol), and 1,2-DCE (7 mL) were combined in a 40 mL vial. The contents were dissolved within several minutes of shaking or stirring. Sodium triacetoxyborohydride (772 mg, 3.64 mmol) was then added as a solid. The reaction was monitored by HPLC and LC-MS and was complete within 3 hours. The reaction was quenched with MeOH (20 mL) and the solvent was subsequently evacuated. The residue was redissolved in 3mL DMF and separated on a C-18 column. Fractions from the prep column dried down in-vacuo and the HCl salt was made by dissolving contents in methanol (sat. HCl). The solvent was reduced and a yellow powder formed. Characterized by ¹H NMR, LC-MS, HPLC.


25

7-Furanyl Sancycline



- 7-iodo sancycline (1.3 mg) and $\text{Pd}(\text{OAc})_2$ were taken in 100 mL of methanol and
5 purged with argon for five minutes at 70 °C. To this solution was added a solution of
sodium carbonate (44 mg) in water (previously purged with argon). A yellow
precipitate was obtained and the mixture was heated for another ten minutes. 3-Furanyl
boronic acid (333 mg, solution in DMF, purged with argon) was then added and the
mixture was heated for another two hours at 70 °C. The reaction was monitored by
10 MPLC/MS. When the reaction was complete, the mixture was filtered through celite
and the solvent was removed to give a crude material. The crude material was purified
by precipitating it with ether (200 ml). The yellow precipitate was filtered and purified
using preparative HPLC. The hydrochloride salt was made by dissolving the material in
MeOH/HCl and evaporating to dryness. The identity of the resulting solid was
15 confirmed using HPLC, MS, and NMR.


7-Cyano Sancycline




- 7-iodo sancycline (1.3 g) was dissolved in NMP (15 mL) and CuCN (344 mg)
20 was added. The reaction mixture was stirred at 80 °C for 15/16 hours overnight. The
reaction mixture was diluted with methanol and centrifuged to yield a grey white
precipitate. The reaction mixture was then passed through Celite and washed with
additional methanol. The filtrate was then concentrated and precipitated with ether. The
solid obtained was then purified using preparative HPLC to yield 7-cyano sancycline in
25 a 50/50 mixture of epimers. The structure of the product was confirmed using mass
spectra and NMR.

7,9-Diiodosancycline

- 30.0 mL of concentrated sulfuric acid was added to 1.00g of sancycline hydrochloride hemihydrate with stirring and the solution cooled to 0°C. 1.09g of N-iodosuccinimide was added portionwise to the solution over one hr and the reaction mixture monitored by HPLC and TLC. The reaction mixture was poured into 250 mL of ice water, extracted three times with n-butanol, and the solvent removed under reduced pressure. The crude residue was purified by preparative HPLC yielding 787mg (61%) of 7-iodosancycline and 291mg (22%) of 7,9-diiodosancycline as yellow and dark yellow crystals respectively.
- MS (FAB): m/z 667 (M+H)
¹H NMR (Methanol d-4, 300MHz) δ 8.35 (s, 1H), 3.78 (s, 1H), 3.33 (s, 2H), 2.88 (s, 7H), 2.41 (m, 2H), 1.41 (m, 5H).

7,9-Bis(3,4-Methylenedioxyphenyl)-Sancycline

- 577 mg (0.74 mmol) 7,9-diiodo sancycline and 8.3 mg (0.37 mmol) palladium acetate were dissolved in 25 ml methanol, under a nitrogen atmosphere. The solution was warmed to 60 °C. After stirring for ten minutes 234 mg (2.22 mmol), sodium carbonate was added followed by 246 mg (1.48 mmol) of 3,4-methylenedioxyphenyl boronic acid. The reaction was complete in 4 hours. The reaction mixture was filtered through a celite bed and concentrated under reduced pressure. This crude product was purified by preparative liquid chromatography using a C₁₈ stationary phase with eluent A: 0.1% TFA in water and eluent B: 0.1% TFA in acetonitrile. 60 mg pure product was isolated.


7-Tetramethylsilylethynyl-Sancycline

SCHEME 10

5

To a solution of 6.54 g (10 mmol) 7-iodo-sancycline trifluoroacetate 500 mg tetrakis-triphenylphosphino-palladate, 500 mg copper(I) iodide, 100 mg palladium acetate and 30 ml triethylamine 3 ml trimethylsilyl-acetylene was added. The reaction mixture was stirred at room temperature for two hours than filtered through a celite bed and concentrated. The dry material was picked up in methanol, the insolubles were filtered out. The solution was concentrated to recover 6.8 g of the product (10B).

10


7-Ethynyl-Sancycline

SCHEME 11

15


7-Tetramethylsilylethynyl-sancycline (10B) is dissolved in 300 ml methanol, and stirred at 40 °C with 6.8 g potassium carbonate. When no starting material could be detected by HPLC (~3 hours), the reaction mixture was cooled in an ice/water bath and solids were removed by filtration. The structure of the alkyne (11B) was confirmed by LCMS. 11B was then used without further purification in the next step.

20

7-Ethyl-Sancycline

SCHEME 12


- 5 10% palladium catalyst on charcoal (1 g) was added to 7-ethynyl sancycline (11B) in a saturated methanol hydrochloric acid solvent. The mixture was placed in a hydrogenator under 50 psi hydrogen pressure. The reaction was completed in ~8 hours. The catalyst was filtered off, and the resulting solution was concentrated. The crude product was purified by preparative liquid chromatography using a C₁₈ stationary phase
- 10 with eluent A: 0.1% TFA in water and eluent B: 0.1% TFA in acetonitrile. The combined clean fractions are concentrated and hydrochloric acid saturated isopropanol added. The pure product is precipitated by addition of diethylether and filtered off. After drying under reduced pressure 3.2 g of 7-ethyl-sancycline (12B) was isolated.

15 7-Ethyl-9-Iodo-Sancycline

SCHEME 13


- 20 7-Ethyl-Sancycline (12B, 6.7 mmol, 3.2 g) was dissolved in 75 ml methanesulfonic acid at room temperature. N-iodo succinimide (13B, 13.5 mmol, 3.05 g) was added over two hours in 6 portions. After two hours diethyl ether was added, and the precipitate was filtered off and dried. The crude product was purified by preparative liquid chromatography using a C₁₈ stationary phase with eluent A: 0.1% TFA in water and eluent B: 0.1% TFA in acetonitrile. 1.5 g of pure product (13C) was isolated.

25

7-Ethyl-9-Cyclohexenylethynyl-Sancycline

SCHEME 14


- 5 To a solution of 7-ethyl-sancycline (500 mg, 1.13 mmol), 50 mg tetrakis-triphenylphosphino-palladate, 50 mg copper(I) iodide, 10 mg palladium acetate and 3 ml triethylamine 0.1 ml cyclohexenyl-acetylene was added. The reaction mixture was stirred at 60 °C for one hour, filtered through a celite bed and concentrated. The dry material was dissolved in methanol and filtered. The solution was then concentrated and
- 10 purified using preparative liquid chromatography. The preparative liquid chromatography used a C₁₈ stationary phase with eluent A: 0.1% TFA in water and eluent B: 0.1% TFA in acetonitrile. 100 mg of the product was isolated.

7-iodo-9-t-Butyl-Sancycline

SCHEME 15

- 9-tbutyl-sancycline (15A, 1.13 g, 2 mmol) was dissolved in 5 ml methanesulfonic acid (0.448, 2 mmol). N-iodosuccinimide (15B) was added at room
- 20 temperature over one hour in four portions. The product (15C) was precipitated with diethyl ether, filtered off and used in other reaction without further purification.


7-(2-Methoxy-5-Dimethylaminomethylphenyl)-9-t-Butyl-Sancycline




SCHEME 16

- 7-Iodo-9-t-butyl-sancycline (15B, 710 mg, 1.0 mmol) and palladium acetate (22.4 mg, 0.1 mmol) were dissolved in 25 ml of methanol under a nitrogen atmosphere. Cesium carbonate (3.25 g, 10 mmol) and 2-methoxy-5-(dimethylaminomethyl)phenylboronic acid (16B, 0.435 g, 0.15 mmol) were added. The reaction mixture was stirred at 60 °C for two hours and then filtered through a celite bed and concentrated under reduced pressure. The crude product was purified by preparative liquid chromatography using a C₁₈ stationary phase with eluent A: 0.1% TFA in water and eluent B: 0.1% TFA in acetonitrile. 210 mg of the product (16C) was isolated.

7-(para-tert-butyl phenyl) -9- aminomethyl sancycline



- 7-para-tert-butyl phenyl sancycline (5.0 g) was dissolved in trifluoroacetic acid (300 mL). Three equivalents of HMBC was added and the reaction was stirred at room temperature. After 72 hours, HPLC indicated that the reaction was complete. The reaction mixture was filtered to give a brown liquid which was subsequently dissolved in methanol and precipitated in diethyl ether. The solid was then purified using HPLC and the product was identified using NMR and mass spectra.

7-Furanyl-9-nitro-Sancycline

500 milligrams of 9-NO₂ sancycline was taken in 20 mL of TFA and cooled down in an ice bath. To this solution, NIS (300 mg) was added in portions and stirred at room temperature for three hours. Once the reaction was completed, 7-iodo-9-NO₂ sancycline was precipitated in diethyl ether. The yellow powder was then filtered and dried *in vacuo*.

7-Iodo-9-nitro-sancycline (585 mg) and PD(OAc)₂ (22 mg) were taken in 20 mL of methanol and purged with argon for five minutes. To this solution, Na₂CO₃ (420 mg, solution in 5 mL H₂O, purged with argon), was added and a yellow precipitate was obtained. The solution was stirred at 55-60 °C for five minutes. To this solution, 3-furanyl boronic acid (160 mg in 5 mL of DMF, purged with argon) was added and the reaction mixture was heated at 70 °C for three hours. The reaction mixture was then passed through celite. Evaporation of the solvent gave a brown solid, which was then recrystallized using a mixture of methanol and ether to yield 7-furanyl 9-nitro sancycline.

7-Furanyl 9-nitro sancycline (500 mg) was taken in 30 ml of methanol. To this solution, PtO₂ (15 mg) was added and hydrogenated at 40 psi for three hours. It was then filtered through celite. The crude material was purified using preparative HPLC to yield 7-furanyl 9-amino sancycline.

Preparation of 9-Iodominocycline

To 200ml of 97% methanesulfonic acid was slowly added, at ambient temperature, portionwise [30g;56.56mM] of minocycline-*bis*-hydrochloride salt. The dark yellow brown solution was then stirred at ambient temperature while [38g;169.7mM] of N-iodosuccinimide was added, in six equal portions, over 3.0 hours time. The reaction was monitored via analytical LC, noting the disappearance of the starting material.

The reaction was slowly quenched into 2L of ice cold water containing [17.88g;1134.1mM] of sodium thiosulfate with rapid stirring. This quench was stirred for approximately 30 minutes at ambient temperature. The aqueous layer was then


- extracted with 6x200ml of ethyl acetate before the aqueous was poured onto [259.8g;3.08M] of sodium hydrogen carbonate containing 300ml of n-butanol. The phases were split and the aqueous extracted with 4x250ml of n-butanol. The organic fractions were combined and washed with 3x250ml of water and once with 250ml of
- 5 saturated brine. The resulting organic phase was reduced to dryness under reduced pressure. The residue was suspended in methanol (~600ml) and anhydrous HCl gas was bubbled into this mixture until solution occurred. This solution was reduced to dryness under reduced pressure. The filtrates were reduced to dryness under reduced pressure. The resulting material was triturated with 300ml of methyl t-butyl ether and isolated via
- 10 filtration. This material was redissolved in 300ml of methanol and treated with 0.5g of wood carbon, filtered and filtrates reduced to dryness under reduced pressure. The material was again powdered under methyl t-butyl ether, isolated via suction filtration and washed with more ether, and finally hexanes. The material was vacuum dried to give 22.6g of a light yellow brown powder.

General Procedure For Preparation of 9-Alkynyl Minocycline Compounds


- 1 mmol 9-iodo minocycline, 50mg tetrakis triphenylphosphinato palladate, 12 mg palladium acetate, 32mg copper (I) iodide are dissolved/suspended in 10ml acetonitrile. 2 to 5ml triethylamine and 3 to 5 mmol alkynyl derivative is added. The reaction mixture is vigorously stirred between ambient temperature to 70°C. The reaction time is 2-24 hours. When the reaction is completed the dark suspension is filtered through a celite bed and concentrated. The crude product is purified by prep HPLC. The combined fractions are concentrated and taken up in ~1ml methanol.. ~3ml HCl saturated methanol is added, and the product is precipitated with ether.

General Procedure For Preparation of 9-Aryl Minocycline Compounds


- 0.15mmol of 9-iodominocycline, PdOAc (3.2mg), 229µl 2M Na₂CO₃ and 2 equivalents of phenyl boronic acid were dissolved/suspended in 10ml methanol. The reaction flask was purged with argon and the reaction run for a minimum of four hours or until HPLC monitoring shows consumption of starting material and/or the appearance of products. The suspension was filtered through celite, and subject to purification by prep HPLC on a divinylbenzene column.

20 9-(4-Trifluoromethoxyphenylureido)-Methyl Minocycline

- To 3 mL of dimethylformamide was added 150 mg (0.25 mmol) of 9-methyl aminominocycline trihydrochloride and 67 mL (0.50 mmol) of triethylamine at 25 °C. With stirring, 75 mL (0.50 mmol) of 4-trifluoromethoxyphenylisocyanate was added and the resulting reaction mixture was stirred at 25 °C for two hours. The reaction was monitored by analytical HPLC (4.6 x 50mm reversed phase Luna C18 column, 5 minute linear gradient 1-100% B buffer, A buffer was water with 0.1% trifluoroacetic acid, B buffer was acetonitrile with 0.1% trifluoroacetic acid). Upon completion, the reaction was quenched with 1 mL of water and the pH adjusted to approximately 2.0 with concentrated HCl. The solution was filtered and the compound purified by preparative HPLC. The yield of the product was 64 mg (37% yield). The purity of the product was 95% determined by LCMS (M+1 = 690).

9-(4'Carboxy phenyl) Minocycline

- In a clean, dry reaction vessel, was placed 9-iodominocycline [500mg; 0.762mmoles]bis HCl salt, palladium (II) acetate [17.2mg; 0.076mmoles] along with 10ml of reagent grade methanol. The solution was immediately purged, with stirring, with a stream of argon gas for approximately 5 minutes. The reaction vessel was brought to reflux and to it was sequentially added via syringe 2M potassium carbonate solution [1.91ml; 3.81mmoles], followed by a solution of p-carboxyphenyl boronic acid [238.3mg; 1.53mmoles]in 5ml of reagent DMF. Both of these solutions were previously degassed with argon gas for approximately 5minutes. The reaction was heated for 45 minutes, the progress was monitored via reverse phase HPLC. The reaction was suctioned filtered through a pad of diatomaceous earth and washed the pad with DMF. The filtrates were reduced to an oil under vacuum and residue treated with t-butylmethyl ether. Crude material was purified via reverse phase HPLC on DVB utilizing a gradient of water and methanol/acetonitrile containing 1.0% trifluoroacetic acid. The product confirmed by mass spectrum: found M+1 578.58; the structure corroborated with ¹H NMR.


20 9-(4'-Acetyl phenyl) Minocycline

- In a clean, dry reaction vessel, was placed 9-iodominocycline (0.762mmoles) bis HCl salt, palladium (II) acetate (0.076mmoles) along with 10ml of reagent grade methanol. The solution was immediately purged, with stirring, with a stream of argon gas for approximately 5 minutes. The reaction vessel was brought to reflux and to it was sequentially added via syringe 2M potassium carbonate solution, followed by a solution of p-acetylphenyl boronic acid (1.53mmoles) in 5ml of reagent DMF. Both of these solutions were previously degassed with argon gas for approximately 5 minutes. The reaction was heated for 45 minutes, the progress was monitored via reverse phase HPLC. The reaction was suctioned filtered through a pad

of diatomaceous earth and washed the pad with DMF. The filtrates were reduced to an oil under vacuum and residue treated with t-butylmethyl ether. Crude material was purified via reverse phase HPLC on DVB utilizing a gradient of water and methanol/acetonitrile containing 1.0% trifluoroacetic acid.

5


N-Benzyl-9'-minocyclinyl guanidine



To a stirred solution of 9-aminominocycline (1.6 mmol) in 30 mL of acetonitrile, benzylcyanamide (6.0 mmol) was added in one portion. The reaction mixture was first heated to reflux at 60 °C for several hours, and continued at room temperature for 4-5 days. The guanidino product was subsequently isolated, and identified using MS, NMR and HPLC.

10

9-Minocycline methyl ester




15

In the Parr apparatus were placed: 9-iodosancycline trifluoroacetic acid salt (0.8 g, 1.17 mmol), NaOAc (0.64g, 4 eq.), Pd(dppf)₂Cl₂, and CH₂Cl₂ (48mg, 5%). The apparatus was closed, purged with CO, and then filled with CO under 450psi. The reaction mixture was stirred for four hours at 80 °C. It was then acidified with TFA and concentrated *in vacuo*. The product was purified by HPLC. A mixture of 3 : 1 epimers was obtained. The yield was 188 mg of product.

20

9-N-piperidinyl-minocycline




25

Concentrated H₂SO₄ (2 mL) was added slowly to a stirred solution of glutaraldehyde (1 mL). Water (0.8 g) was added and stirred at room temperature for


- eighteen hours and heater to 70 °C for two hours. The mixture was then cooled to room temperature. The solution was then transferred to a solution of 9-amino minocycline in DMF (5 ml) and stirred at room temperature for two days until all starting material was consumed, as indicated by HPLC. The product was isolated and purified using standard techniques. The structure of the product was confirmed by NMR and mass spec.

2-[4-(5-Minocyclin-9-yl-furan-2-ylmethyl)-piperazin-1-yl]-ethanol



- Na_2CO_3 (0.64 g) in water (5 mL) was added to a degassed solution of 9-iodo-minocycline hydrochloride (1 g) and $\text{Pd}(\text{OAc})_2$ (100 mg) in methanol (10mL). The reaction was stirred for five minutes at 60 °C. 2-Formyl furan-5-boronic acid (0.3 g) in methanol (10 mL) was then added, and the reaction was allowed to proceed for four hours. The mixture was then filtered and concentrated to give a brown solid (9-(2'formyl furanyl)-minocycline).
- The brown solid (9-(2'formyl furanyl)-minocycline, 1 g) was dissolved in 20 mL of methanol and acetic acid (2 mL) and hydroxyethyl piperazine (1 mL) was added and stirred for ten minutes at room temperature. The reaction was quenched with ether (200 mL), and the organic layer was then washed and concentrated to yield a brown oil. The brown oil was the dissolved in methanol (10 mL) and water. The mixture was the chromatographed using a CH_3CN gradient to yield the product, 2-[4-(9-Minocyclin-2-yl-furan-2-ylmethyl)-piperazin-1-yl]-ethanol. The product was confirmed using MS, NMR, and HPLC.

9-N-morpholinyl minocycline




25

NaCNBH_3 (200 mg) was added to a stirred solution of 9-amino minocycline H_2SO_4 (1 g) in methanol (4.9 mL) and acetic acid 91 mL) and stirred for five minutes at room temperature. (2-Oxo-ethoxy)-acetaldehyde (10 mL) was added dropwise and stirred for fifteen minutes at room temperature. The reaction mixture was

concentrated with out heat and the residue was dissolved in 20 mL of methanol and TFA (0.5 mL). The product was obtained using preparative HPLC and converted to the HCl salt. The product was confirmed using mass spectra and NMR.

5 N-Benzyl-N',N'-dimethyl-N-(5-minocyclin-9-yl-furan-2-ylmethyl)-ethane-1,2-diamine



Na_2CO_3 (0.64 g) in water (5 mL) was added to a degassed solution of 9-iodo-minocycline hydrochloride (1 g) and $\text{Pd}(\text{OAc})_2$ (100 mg) in methanol (10mL). The reaction was stirred for five minutes at 60 °C. 2-Formyl furan-5-boronic acid (0.3 g) in
10 methanol (10 mL) was then added, and the reaction was allowed to proceed for four hours. The mixture was then filtered and concentrated to give a brown solid (9-(2'-formyl furanyl)-minocycline).

The brown solid (9-(2'-formyl furanyl)-minocycline, 1 g) was dissolved in 20 mL of methanol and acetic acid (2 mL) and N'-benzyl-N,N-dimethyl
15 ethylenediamine (1 mL) was added and stirred for ten minutes at room temperature. The reaction was quenched with ether (200 mL), and the organic layer was then washed and concentrated to yield a brown oil. The brown oil was the dissolved in methanol (10 mL) and water. The mixture was the chromatographed using a CH_3CN gradient to yield the
20 product, N-Benzyl-N',N'-dimethyl-N-(5-minocyclin-9-yl-furan-2-ylmethyl)-ethane-1,2-diamine. The product was confirmed using MS, NMR, and HPLC.

[4S-(4 α ,12 α)]-9-Amino-4-dimethylamino-3,5,10,12,12a-pentahydroxy-6-methyl-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydro-naphthacene-2-carboxamide

9-Nitro doxycycline hydrochloride was synthesized as previously
25 described. 1.0 g was hydrogenated in 40 mL MeOH, 1 mL conc. HCl and 100 mg of 10% Pd/C for 3 hr under 30 psi of H_2 . The solution was filtered through Celite and the filtrate evaporated *in vacuo* to obtain 0.9 g of the dihydrochloride salt as a yellow solid. Yield = 82%. ^1H NMR (CD_3OD , 300 MHz): δ 7.62 (d, 1H, J = 8.08 Hz); 7.14 (d, 1H, J = 8.08); 4.42 (s, 1H); 3.6 (dd, 1H); 2.98, 2.90 (each s, 3H); 2.84 (d, 1H); 2.72 (m, 1H);
30 2.59 (dd, 1H); 1.56 (d, 3H). HRMS calcd ($\text{C}_{22}\text{H}_{26}\text{N}_3\text{O}_8+\text{H}$) 460.1720, found 460.1739.

[4S-(4 α ,12 α)]-9-(Diazonium)-4-dimethylamino-3,5,10,12,12a-pentahydroxy-6-methyl-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydro-naphthacene-2-carboxamide

A flask was charged with 100 mg of compound 10 and dissolved with 4 mL of 0.1 N HCl. The solution was cooled to 0 ° C and 35 μ L of butyl nitrite was added with stirring. After 1 h, the bright red mixture was added dropwise to 100 mL of cold anhydrous Et₂O. The product was collected by filtration under ether as an orange solid. Analytical R_t = 1.90 Method A. LC/MS(ESI): 472 (M+H).

10 *General Diazonium Reaction Procedure*

0.1 g of 9-diazonium salt generated *in situ* (HCl salt) or tetrafluoroborate salt was dissolved in MeOH and 0.05 equiv. of Pd(OAc)₂ added and up to 10% acetic acid. The reaction mixture was stirred for 5 minutes at room temperature, and 2 or more equiv. of the desired reactant was added. The reaction was typically continued for 18 hours. The catalyst was removed and the filtrate treated with activated charcoal and dried to give the crude product.

[4S-(4 α , 12 α)]-9-(3-Oxo-propenyl)-4-dimethylamino-3,5,10,12,12a-pentahydroxy-6-methyl-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydro-naphthacene-2-carboxamide

Preparative C18 reverse-phase HPLC. R_t = 17.3. Analytical R_t = 9.42. ¹H NMR (CD₃OD, 300 MHz): δ 7.69 (d, 1H, *J* = 18.0 Hz); 7.70 (d, 1H, *J* = 7.8 Hz); 7.32 (d, 1H, *J* = 7.8); 6.65 (d, 1H, *J* = 18.0 Hz); 4.05 (s, 1H); 3.57 (dd, 1H); 2.94 (s, 6H); 2.85 (d, 1H); 2.71 (m, 1H); 2.60 (dd, 1H); 1.55 (d, 3H). MS(FAB): 499 (M+H).

25 [4S-(4 α , 12 α)]-9-[1'(*E*)-(2'-Phenyl)ethenyl]-4-dimethylamino-3,5,10,12,12a-pentahydroxy-6-methyl-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydro-naphthacene-2-carboxamide

Preparative C18 reverse-phase HPLC R_t = 27-30. Analytical R_t = 19.42. ¹H NMR (CD₃OD, 300 MHz): δ 7.69 (d, 1H, *J* = 8.0 Hz); 7.37 (d, 1H, *J* = 24.0 Hz); 7.14 (m, 6H); 6.74 (d, 1H, *J* = 8.0); 4.04 (s, 1H); 3.49 (d, 1H); 2.91 (s, 6H); 2.8 (d, 1H); 2.70 (m, 1H); 2.55 (d, 1H); 1.36 (m, 3H). MS(FAB): 547.1. HRMS calcd (C₃₀H₃₀N₂O₈+H) 547.2082, found 547.2080.

35


General Aryl or Heteroaryl Boronic Acids Reaction Procedure

A solution of diazonium salt (HCl or HBF₄) 14 in MeOH (approx. 10 mg/mL) was cooled to 0 °C and 0.1 equiv. of Pd(OAc)₂ was added. The mixture was stirred for 5 min, and 1 or more equiv. of phenylboronic acid or heteroaryl boronic was added and stirred for 6 h, warming to room temperature during the reaction. The catalyst was filtered away through Celite, and the filtrate dried down to yield the crude reaction product which was further purified by C18 reverse-phase preparative chromatography.

10 [4S-(4 α ,12 α)]-9-(4-Methoxy-phenyl)-4-dimethylamino-3,5,10,12,12a-pentahydroxy-6-methyl-1,11-dioxo-1,4,4a,5,5a,6,11,12a-octahydro-naphthacene-2-carboxamide

The purified product was obtained by preparative C18 reverse-phase HPLC, R_f = 33.97. Analytical R_f = 18.2. ¹H NMR (CD₃ OD, 300 MHz): δ = 7.50 (m, 3H); 6.94 (m, 3H); 4.16 (m, 1H); 3.82 (d, 1H); 3.30 (s, 3H); 2.79 (s, 6H); 2.74 (m, 1H); 2.56 (m, 1H); 1.53 (d, 3H); MS(FAB): 551 (M+H).

9-Carbamoyl-4-chloro-7-dimethylamino-1,8,10a,11-tetrahydroxy-10,12-dioxo-6,6a,7,10,10a,12-hexahydro-naphthacene-2-sulfonic acid



20

Trifluoroacetic Acid (30 mL), and 3ml of sulfuric acid were taken in a 100ml flask and stirred for five minutes. 7-chloro sancycline (4.3 mmol) was added to the acid mixture and stirred for 2-12 h at room temperature to 50 degree. After completion (monitored by LC/MS), the mixture was precipitated in 300ml ether, filtered and dried. The product was then purified using preparative HPLC.

25

¹H NMR δ 7.95 (1H, s), 7.15 (1H, s), 4.22 (1H, s), 3.65-3.59 (2H, m), 3.15 (6H, s), 3.09 (1H, m), 2.99 (1H, s), 2.86 (1H, s). M+1 527.18. CHN Analysis : Calc. C 43.10, H 3.15, N 4.37, S 5.00; found C 42.26, H 3.73, N 4.50, S 5.31

30

Example 2: *In vitro* Minimum Inhibitory Concentration (MIC) Assay

The following assay is used to determine the efficacy of the tetracycline compounds against common bacteria. 2 mg of each compound is dissolved in 100 μ l of DMSO. The solution is then added to cation-adjusted Mueller Hinton broth (CAMHB), which results in a final compound concentration of 200 μ g per ml. The tetracycline compound solutions are diluted to 50 μ L volumes, with a test compound concentration of .098 μ g/ml. Optical density (OD) determinations are made from fresh log-phase broth cultures of the test strains. Dilutions are made to achieve a final cell density of 1×10^6 CFU/ml. At OD=1, cell densities for different genera should be approximately:

10

<i>E. coli</i>	1×10^9 CFU/ml
<i>S. aureus</i>	5×10^8 CFU/ml
<i>Enterococcus sp.</i>	2.5×10^9 CFU/ml

15

50 μ l of the cell suspensions are added to each well of microtiter plates. The final cell density should be approximately 5×10^5 CFU/ml. These plates are incubated at 35°C in an ambient air incubator for approximately 18 hr. The plates are read with a microplate reader and are visually inspected when necessary. The MIC is defined as the lowest concentration of the tetracycline compound that inhibits growth.

20

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described herein. Such equivalents are considered to be within the scope of the present invention and are covered by the following claims. The contents of all references, patents, and patent applications cited throughout this application are hereby incorporated by reference. The appropriate components, processes, and methods of those patents, applications and other documents may be selected for the present invention and embodiments thereof.

30

This application is related to U.S. Patent Application Serial No. 09/895,812, entitled "7- Substituted Tetracycline Compounds," filed June 29, 2001; U.S. Provisional Patent Application Serial No. 60/275,576, entitled "7-Substituted Tetracycline Compounds" filed March 13, 2001; and U.S. Provisional Patent Application Serial No. 60/216,760, entitled "7-Substituted Sancycline Compounds" filed on July 7, 2000; the entire contents of each of these applications are hereby incorporated herein by reference.


35

This application is related to U.S. Patent Application Serial No. 09/895,797, entitled "7,9-Substituted Tetracycline Compounds," filed on June 29, 2001; and U.S. Provisional Patent Application Serial No. 60/275,620, entitled "7, 9-Substituted Tetracycline Compounds," filed on March 13, 2001. The entire contents of
5 each of these applications are incorporated herein by reference.

This application is related to U.S. Patent Application Serial No. 09/895,857, entitled "9 Substituted Minocycline Compounds," filed on June 29, 2001; U.S. Provisional Patent Application Serial No. 60/275,621, entitled "9-Substituted Minocycline Compounds," filed on March 13, 2001; and U.S. Provisional Patent
10 Application Serial No. 60/216,659, entitled "9-Substituted Minocycline Compounds," filed on July 7, 2000; each of these applications are hereby incorporated herein by reference.

CLAIMS

1. A substituted tetracycline compound of Formula I:



5


wherein:

- X is $\text{CHC}(\text{R}^{13}\text{Y}^*\text{Y})$, CR^6R^6 , $\text{C}=\text{CR}^6\text{R}^6$, S, NR^6 , or O;
 R^2 , $\text{R}^{2'}$, R^4 , and $\text{R}^{4''}$ are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;
 R^4 is $\text{NR}^4\text{R}^{4''}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;
 $\text{R}^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;
 R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;
 R^6 and R^6 are each independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
 R^7 is nitro, heterocyclic, alkyl, aminoalkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, carbonyl, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, or $-(\text{CH}_2)_{0-3}\text{NR}^{7c}\text{C}(=\text{W}')\text{WR}^{7a}$;
 R^8 is hydrogen, hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(\text{CH}_2)_{0-3}\text{NR}^{8c}\text{C}(=\text{E}')\text{ER}^{8a}$;
 R^9 is hydrogen, hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(\text{CH}_2)_{0-3}\text{NR}^{9c}\text{C}(=\text{Z}')\text{ZR}^{9a}$;
 R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} , R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} , R^{8f} , R^{9a} , R^{9b} , R^{9c} , R^{9d} , R^{9e} , and R^{8f} are each independently absent, hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

- R¹³ is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, aryl, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
 E is CR^{8d}R^{8e}, S, NR^{8b} or O;
 E' is O, NR^{8f}, or S;
 5 Z is CR^{9d}R^{9e}, S, NR^{9b} or O;
 Z' is O, S, or NR^{9f};
 W is CR^{7d}R^{7e}, S, NR^{7b} or O;
 W' is O, NR^{7f} S;
 R¹³ is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio,
 10 alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl; and
 Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts thereof.
- 15 2. The tetracycline compound of claim 1, wherein R⁴ is NR^{4'}R^{4''}, X is CR⁶R^{6'}; R², R^{2'}, R⁶, R^{6'}, R⁸, R⁹, R¹⁰, R¹¹, and R¹² are each hydrogen; R^{4'} and R^{4''} are lower alkyl; and R⁵ is hydroxy or hydrogen.
3. The tetracycline compound of claim 2, wherein R^{4'} and R^{4''} are each
 20 methyl and R⁵ is hydrogen.
- 4 The tetracycline compound of claim 3 wherein R⁷ is furanyl.
5. The tetracycline compound of claim 4, wherein said furanyl is substituted
 25 with a substituted or unsubstituted alkyl group.
6. The tetracycline compound of claim 5, wherein said furanyl group is substituted with an alkyl group further substituted with one or more heterocyclic groups.
- 30 7. The tetracycline compound of claim 3, wherein R⁷ is morpholinyl, piperidinyl, or piperazinyl.
8. The tetracycline compound of claim 3, wherein R⁷ is of the formula: -
 (C=O)-R^{7j}, wherein R^{7j} is alkcarbonyl, alkyl, alkenyl, alkynyl, aryl, heterocyclic or
 35 aralkyl.

9. The tetracycline compound of claim 8, wherein R^{7j} is substituted or unsubstituted alkyl.
10. The tetracycline compound of claim 9, wherein R^{7j} is alkyl and
5 substituted with a heterocycle.
11. The tetracycline compound of claim 10, wherein said heterocycle is morpholinyl, piperidinyl, or piperazinyl.
- 10 12. The tetracycline compound of claim 3, wherein R^7 is phenyl substituted with sulfonyl amino or alkylcarbonylamino.
13. The tetracycline compound of claim 3, wherein R^7 is substituted pyrrole.
- 15 14. The tetracycline compound of claim 13, wherein said pyrrole is substituted with a carbonyl moiety.
15. The tetracycline compound of claim 14, wherein said carbonyl moiety is alkoxycarbonyl.
20
16. The tetracycline compound of claim 3, wherein R^7 is substituted pyrazine.
17. The tetracycline compound of claim 16, wherein said substituted
25 pyrazine is substituted with an substituted or unsubstituted alkoxycarbonylamino group, an alkaminocarbonylamino group, or an alkylaminocarbonyl group.
18. The tetracycline compound of claim 3, wherein R^7 is $-C(=W')WR^{7a}$, wherein W' is NR^{7f} , W is $CR^{7d}R^{7e}$, and R^{7a} is hydrogen.
30
19. The tetracycline compound of claim 18, wherein R^{7f} is alkoxy.
20. The tetracycline compound of claim 3, wherein R^7 is $-NR^{7c}C(=W')WR^{7a}$, wherein R^{7c} is hydrogen, W' is NH , W is NR^{7b} , and R^{7a} and R^{7b} taken together are
35 heterocyclic.

21. The tetracycline compound of claim 3, wherein R^7 is a boronic acid.
22. The tetracycline compound of claim 21, wherein R^7 is $-B(OR^{b1})(OR^{b2})$, wherein R^{b1} and R^{b2} are each hydrogen or alkyl, and optionally linked to form a ring.
- 5 23. The tetracycline compound of claim 3, wherein R^7 is a sulfonic acid.
24. The tetracycline compound of claim 23, wherein R^7 is $-SO_3H$.
- 10 25. The tetracycline compound of claim 3, wherein R^7 is isoxazolyl.
26. A substituted tetracycline compound of Formula II:



(II)

15 wherein:

- X is $CHC(R^{13}Y'Y)$, CR^6R^6 , S, NR^6 , or O;
- R^2 , R^2 , R^4 , and R^4 are each independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;
- 20 R^4 is NR^4R^4 , alkyl, alkenyl, alkynyl, hydroxyl, halogen, or hydrogen;
- R^2 , R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;
- R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;
- 25 R^6 and R^6 are each independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
- R^7 is $-CH_2NR^{7a}R^{7b}$, halogen, $-CH_2OR^{7a}$, substituted alkenylfuranyl, pyrazinyl, pyridinyl, alkyl, acyl, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, or $-(CH_2)_0$;
- 30 $NR^{7c}C(=W')WR^{7a}$;

R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^9 is $-\text{CH}_2\text{SR}^{9a}$, $-\text{CH}_2\text{S}(=\text{O})\text{R}^{9a}$, $-\text{CH}_2\text{S}(=\text{O})_2\text{R}^{9a}$, $-\text{CH}_2\text{NR}^{9a}\text{R}^{9b}$, SO_3H , aminoalkyl, furanyl, substituted alkyl, $-(\text{CH}_2)_{0-3}(\text{NR}^{9c})_{0-1}\text{C}(=\text{Z}')(\text{Z})_{0-1}\text{R}^{9a}$, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, thionitroso, or $-(\text{CH}_2)_{0-3}(\text{NR}^{9c})_{0-1}\text{C}(=\text{Z}')(\text{Z})_{0-1}\text{R}^{9a}$;

Z is $\text{CR}^{9d}\text{R}^{9e}$, S , NR^{9b} or O ;

Z' is O , S , or NR^{9f} ;

W is $\text{CR}^{7d}\text{R}^{7e}$, S , NR^{7b} or O ;

W' is O , NR^{7f} S ;

R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} , R^{9a} , R^{9b} , R^{9c} , R^{9d} , and R^{9e} are each independently hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl; and

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts thereof, provided that R^7 and R^9 are not both unsubstituted phenyl.

20

27. The tetracycline compound of claim 26, wherein X is CR^6R^6 ; R^2 , R^2 , R^6 , R^6 , R^8 , R^{10} , R^{11} , and R^{12} are each hydrogen; R^4 is $\text{NR}^{4'}\text{R}^{4''}$; $R^{4'}$ and $R^{4''}$ are lower alkyl; and R^5 is hydroxy or hydrogen.

25 28. The tetracycline compound of claim 27, wherein $R^{4'}$ and $R^{4''}$ are each methyl and R^5 is hydrogen.

29. The tetracycline compound of claim 26, wherein R^7 is furanyl.

30 30. The tetracycline compound of claim 26, wherein R^7 is ethyl.

31. The tetracycline compound of claim 26, wherein R^7 is $-\text{CH}_2\text{O-alkyl}$ and R^9 is alkyl.


35 32. The tetracycline compound of claim 26, wherein R^7 is pyridinyl and R^9 is alkyl substituted with piperidine or alkylamino.

33. The tetracycline compound of claim 26, wherein R^7 is iodo or chloro.

34. The tetracycline compound of claim 26, wherein R^9 is alkyl, $-\text{CH}_2\text{-S-R}^{9h}$,
 5 $-\text{CH}_2\text{-S(=O)-alkyl}$, $-\text{CH}_2\text{-S(=O)}_2\text{-alkyl}$, $-\text{SO}_3\text{H}$, amino, nitro, or $-\text{CH}_2\text{O-alkyl}$.

35. The tetracycline compound of claim 19, wherein R^{9h} is is alkyl, alkenyl, alkynyl, aryl, aralkyl, heterocyclic or hydrogen.

10 36. A tetracycline compound of formula III:



wherein:

X is $\text{CHC(R}^{13}\text{Y'Y)}$, $\text{CR}^{6'}\text{R}^6$, S, NR^6 , or O;

15 R^2 , $R^{4'}$, $R^{4''}$, $R^{7'}$ and $R^{7''}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R^4 is $\text{NR}^{4'}\text{R}^{4''}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;

$R^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

20 R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;

R^6 and $R^{6'}$ are independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl,
 25 alkylamino, or an arylalkyl;

R^7 is $\text{NR}^{7'}\text{R}^{7''}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;

R^8 is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^9 is $-\text{C(=Z')R}^{9a}$, $\text{CH}_2\text{S(=O)R}^{9a}$, $-\text{CH}_2\text{OR}^{9a}$, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, heterocyclic, arylalkenyl, arylalkynyl, thionitroso, substituted
 30 alkyl, or $-(\text{CH}_2)_{0-3}(\text{NR}^{9c})_{0-1}\text{C(=Z')}(Z)_{0-1}\text{R}^{9a}$;

Z is $\text{CR}^{9d}\text{R}^{9e}$, S, NR^{9b} or O;

Z' is NR^{9f}, O or S;

R^{9a}, R^{9b}, R^{9c}, R^{9d}, R^{9e} and R^{9f} are each independently hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

- 5 R⁸ is hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R¹³ is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

- 10 Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts, esters and prodrugs thereof.

37. The tetracycline compound of claim 36, wherein R⁴ is NR^{4'}R^{4''}; X is
15 CR⁶R^{6'}; R⁷ is NR^{7'}R^{7''}, R², R^{2'}, R⁵, R⁶, R^{6'}, R⁸, R⁹, R¹⁰, R¹¹, and R¹² are each hydrogen; and, R^{4'}, R^{4''}, R^{7'}, and R^{7''} are each lower alkyl.

38. The tetracycline compound of claim 37, wherein R^{4'}, R^{4''}, R^{7'}, and R^{7''}
20 are each methyl.

39. The tetracycline compound of claim 37, wherein R⁷ is hydrogen.

40. The tetracycline compound of claim 36, wherein R⁹ is amino substituted
25 with morpholinyl, piperazinyl, or piperidinyl.

41. The tetracycline compound of claim 36, wherein R⁹ is furanyl.

42. The tetracycline compound of claim 41, wherein said furanyl is
substituted with one or more heterocyclic groups.


- 30 43. The tetracycline compound of claim 36, wherein R⁹ is -C(=Z')R^{9a}, wherein Z' is NR^{9f}, and R^{9a} is hydrogen.

44. The tetracycline compound of claim 43, wherein R^{9f} is alkoxy.
35

45. The tetracycline compound of claim 36, wherein said heterocycle is 2,3-dihydro-isoinidole.

46. A tetracycline compound of formula IV:

5



wherein:

- X is $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$, CR^6R^6 , S, NR^6 , or O;
 R^2 , R^4 , $\text{R}^{4'}$, $\text{R}^{7'}$ and $\text{R}^{7''}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy,
 10 alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;
 R^4 is $\text{NR}^{4'}\text{R}^{4''}$, alkyl, alkenyl, alkynyl, aryl, hydroxyl, halogen, or hydrogen;
 $\text{R}^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;
 R^5 is hydroxyl, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic,
 15 alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, alkyl carbonyloxy, or aryl carbonyloxy;
 R^6 and $\text{R}^{6'}$ are independently hydrogen, methylene, absent, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;
 20 R^7 is hydrogen, hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(\text{CH}_2)_{0-3}(\text{NR}^{7c})_{0-1}\text{C}(=\text{W}')\text{WR}^{7a}$;
 R^8 is hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, amino, arylalkenyl, arylalkynyl, acyl,
 25 aminoalkyl, heterocyclic, thionitroso, or $-(\text{CH}_2)_{0-3}(\text{NR}^{8c})_{0-1}\text{C}(=\text{E}')\text{ER}^{8a}$;
 R^9 is hydrogen, hydroxyl, halogen, thiol, nitro, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, arylalkyl, amino, arylalkenyl, arylalkynyl, acyl, aminoalkyl, heterocyclic, thionitroso, or $-(\text{CH}_2)_{0-3}\text{NR}^{9c}\text{C}(=\text{Z}')\text{ZR}^{9a}$;
 R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} , R^{7f} , R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} , R^{8f} , R^{9a} , R^{9b} , R^{9c} , R^{9d} , R^{9e} , and
 30 R^{8f} are each independently absent, hydrogen, acyl, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

E is $\text{CR}^{8d}\text{R}^{8e}$, S, NR^{8b} or O;

E' is O, NR^{8f} , or S;

W is $\text{CR}^{7d}\text{R}^{7e}$, S, O or NR^{7b} ;

W' is O, NR^{7f} , or S;

- 5 R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl, and pharmaceutically acceptable salts, esters and prodrugs

10 thereof.

47. The tetracycline compound of claim 46, wherein X is CR^6R^6 ; R^2 , $\text{R}^{2'}$, R^6 , $\text{R}^{6'}$, R^8 , R^{10} , R^{11} , and R^{12} are each hydrogen; R^4 is $\text{NR}^{4'}\text{R}^{4''}$; $\text{R}^{4'}$ and $\text{R}^{4''}$ are lower alkyl; and R^5 is hydroxy or hydrogen.

15

48. The tetracycline compound of claim 47, wherein R^8 is halogen.

49. The tetracycline compound of claim 47, wherein R^8 is aryl.




20 50. The tetracycline compound of claim 49, wherein R^8 is furanyl.



















51. The tetracycline compound of claim 49, wherein R^8 is phenyl.


52. The tetracycline compound of claim 48 or 49, wherein R^9 is substituted
25 or unsubstituted amino.


53. The tetracycline compound of claim 48 or 49, wherein R^7 is substituted or unsubstituted amino.
















30 54. A tetracycline compound selected from the group consisting of:






























































































and pharmaceutically acceptable esters, enantiomers, amides, prodrugs, and salts thereof.

55. A method for treating a tetracycline responsive state in a subject,
- 5 comprising administering to said subject a tetracycline compound of any one of claims 1, 26, 36, 46, or 54, such that said subject is treated.
56. The method of claim 55, wherein said tetracycline responsive state is a bacterial infection.
- 10 57. The method of claim 56, wherein said bacterial infection is associated with *E. coli*.
58. The method of claim 56, wherein said bacterial infection is associated
- 15 with *S. aureus*.
59. The method of claim 56, wherein said bacterial infection is associated with *E. faecalis*.
- 20 60. The method of claim 56, wherein said bacterial infection is resistant to other tetracycline antibiotics.
61. The method of claim 55, wherein said subject is a human.
- 25 62. The method of claim 55, wherein said tetracycline compound is administered with a pharmaceutically acceptable carrier.

63. A pharmaceutical composition comprising a therapeutically effective amount of a tetracycline compound of any one of claims 1, 26, 36, 46, or 54 and a pharmaceutically acceptable carrier.

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 October 2003 (02.10.2003)

PCT

(10) International Publication Number
WO 2003/079984 A3

(51) International Patent Classification⁷: C07C 237/26,
43/00, 49/00

(21) International Application Number:
PCT/US2003/008324

(22) International Filing Date: 18 March 2003 (18.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/367,045 21 March 2002 (21.03.2002) US
60/366,915 21 March 2002 (21.03.2002) US
60/367,048 21 March 2002 (21.03.2002) US
60/395,468 12 July 2002 (12.07.2002) US
60/440,305 14 January 2003 (14.01.2003) US

(71) Applicant: PARATEK PHARMACEUTICALS, INC.
[US/US]; 75 Kneeland Street, Boston, MA 02111 (US).

(72) Inventors: NELSON, Mark, L.; 735 Worcester Road, Wellesley, MA 02481 (US). OHMENG, Kwasi; 400 Engamore Lane, Apt. 101, Norwood, MA 02062 (US). FRECHETTE, Roger; 40 Estate Lane, Reading, MA 01867 (US). ABATO, Paul; 41 Pembroke Avenue, Providence, RI 02408 (US). ASSEFA, Haregewein; 29 McCusker Drive, #5, Braintree, MA 02184 (US). BANDARAGE, Upul; 11 Bates Road, Lexington, MA 02421 (US). BERNIAC, Joel; 21 Sunrise Avenue, Stoneham, MA 02180 (US). BHATIA, Beena; 3 Brian Lane, Mansfield, MA 02048 (US). CHEN, Jackson; 1731 Beacon Street, #204, Brookline, MA 02445 (US). ISMAIL, Mohamed, Y.; 5 Hunt Road, Bedford, MA 01730 (US). KIM, Oak, A.; 86 Ellery Street, #1 Apt. 24, Cambridge, MA 02138 (US). MATHEWS, Jude; 633 Moody Street, Waltham, MA 02453 (US). MCINTYRE, Laura; 27 Thorndike Street, Arlington, MA 02474 (US). NIHLAWI, Mohammed; 230 Willard Street, #512, Quincy, MA 02169 (US). PEARSON, Andre; 23 Cutter Street, 32, Somerville, MA 02143 (US). REDDY, Laxma; 49 Eldred

Street, Lexington, MA 02420 (US). SHEAHAN, Paul; 4174 Bannock Avenue, San Diego, CA 92117 (US). SIZENSKY, Emmanuelle; 4 Blackstone Lane, South Grafton, MA 01560 (US). TOURIGNY, Justin; 42 Brainerd Road, #1, Allston, MA 02134 (US). VERMA, Atul, K.; 3 Brian Lane, Mansfield, MA 02048 (US). VISKI, Peter; 887 West Roxbury Parkway, Brookline, MA 02467 (US). WARCHOL, Tadeusz; 177 South Street, Northborough, MA 01532 (US).

(74) Agents: HANLEY, Elizabeth, A. et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
10 September 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED TETRACYCLINE COMPOUNDS

(57) Abstract: The present invention pertains, at least in part, to novel substituted tetracycline compounds. These tetracycline compounds can be used to treat numerous tetracycline compound-responsive states, such as bacterial infections and neoplasms, as well as other known applications for minocycline and tetracycline compounds in general, such as blocking tetracycline efflux and modulation of gene expression.

WO 2003/079984 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/08324

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07C 237/26, 43/00, 49/00

US CL : 552/203, 205, 206; 514/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 552/203, 205, 206; 514/16

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

USPATFULL, STN, HCAPLUS, REGISTRY

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5495018 A (SUM et al) 27 February 1996 (27.02.1996), see the entire document, especially the examples and claims.	1-63
Y	Database CAPLUS on ACS, Accession No: 1994: 217052, Glycylcyclines. A new generation of potent antibacterial agents through modification of 9-aminotetracyclines. Journal of Medicinal Chemistry. Sum, P-E et al; 1994, vol. 37, no.1, pages 184-8, see abstract.	1-63

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 December 2003 (13.12.2003)

Date of mailing of the international search report

03 AUG 2004

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US

Commissioner for Patents

P.O. Box 1450

Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Sabih Qazi

Telephone No: 703-308-1235